

μ -CONSTANT MONODROMY GROUPS AND TORELLI RESULTS FOR MARKED SINGULARITIES, FOR THE UNIMODAL AND SOME BIMODAL SINGULARITIES

FALKO GAUSS AND CLAUS HERTLING

ABSTRACT. This paper is a sequel to [He7]. There a notion of marking of isolated hypersurface singularities was defined, and a moduli space M_μ^{mar} for marked singularities in one μ -homotopy class of isolated hypersurface singularities was established. One can consider it as a global μ -constant stratum or as a Teichmüller space for singularities. It comes together with a μ -constant monodromy group $G^{mar} \subset G_{\mathbb{Z}}$. Here $G_{\mathbb{Z}}$ is the group of automorphisms of a Milnor lattice which respect the Seifert form. It was conjectured that M_μ^{mar} is connected. This is equivalent to $G^{mar} = G_{\mathbb{Z}}$. Also Torelli type conjectures were formulated. All conjectures were proved for the simple singularities and 22 of the exceptional unimodal and bimodal singularities. In this paper the conjectures are proved for the remaining unimodal singularities and the remaining exceptional bimodal singularities.

CONTENTS

1. Introduction	2
2. Review on the topology of isolated hypersurface singularities	4
3. The group $G_{\mathbb{Z}}$ for the simple elliptic and the hyperbolic singularities	9
4. The group $G_{\mathbb{Z}}$ for 6 of the 28 exceptional unimodal and bimodal singularities	15
5. Review on μ -constant monodromy groups G^{mar} , marked singularities, their moduli spaces M_μ^{mar} , and Torelli type conjectures	21
6. G^{mar} , M_μ^{mar} and a strong Torelli result for the simple elliptic and the hyperbolic singularities	28

Date: April 11, 2016.

2000 Mathematics Subject Classification. 32S15, 32S40, 14D22, 58K70.

Key words and phrases. μ -constant monodromy group, marked singularity, moduli space, Torelli type problem, simple elliptic singularities, hyperbolic singularities.

This work was supported by the DFG grant He2287/4-1 (SISYPH).

7. G^{mar} , M_μ^{mar} and a strong Torelli result for 6 of the 28 exceptional unimodal and bimodal singularities	31
8. More on $G_\mathbb{Z}$ for the simple elliptic singularities	34
References	38

1. INTRODUCTION

This paper is a sequel to [He7]. That paper studied local objects, namely holomorphic functions germs $f : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$ with an isolated singularity at 0 (short: singularity), from a global perspective.

There a notion of marking of a singularity was defined. One has to fix one singularity f_0 , which serves as reference singularity. Then a marked singularity is a pair $(f, \pm\rho)$ where f is in the μ -homotopy class of f_0 and $\rho : (Ml(f), L) \rightarrow (Ml(f_0), L)$ is an isomorphism. Here $Ml(f)$ is the Milnor lattice of f , and L is the Seifert form on it (the definitions are recalled in section 2). The group $G_\mathbb{Z}(f_0) := \text{Aut}(Ml(f_0), L)$ will be important, too.

A moduli space M_μ^{mar} of right equivalence classes of marked singularities in one μ -homotopy class was established. One can consider it as a global μ -constant stratum or as a Teichmüller space for singularities. The group $G_\mathbb{Z}(f_0)$ acts properly discontinuously on it, and the quotient is the moduli space M_μ of right equivalence classes of unmarked singularities from [He6, chapter 13]. The μ -constant monodromy group $G^{mar}(f_0) \subset G_\mathbb{Z}(f_0)$ is the subgroup of automorphisms which map the topological component $(M_\mu^{mar}(f_0))^0$ which contains $[(f_0, \pm \text{id})]$ to itself. It can also be constructed as the group of all automorphisms which can be realized modulo $\pm \text{id}$ as transversal monodromies of μ -constant families.

Conjecture 1.1. [He7, conjecture 3.2 (a)] M_μ^{mar} is connected. Equivalently: $G^{mar} = G_\mathbb{Z}$.

Roughly this conjecture says that all abstract automorphisms come from geometry, from coordinate changes and μ -constant families.

Also Torelli type conjectures are formulated in [He7]. Any singularity f comes equipped with its *Brieskorn lattice* $H_0''(f)$, and any marked singularity $(f, \pm\rho)$ comes equipped with a *marked Brieskorn lattice* $BL(f, \pm\rho)$. The Gauß-Manin connection for singularities and the Brieskorn lattices had been introduced 1970 by Brieskorn and had been studied since then. The second author has a long-going project on Torelli type conjectures around them.

In [He4] a classifying space $D_{BL}(f_0)$ for marked Brieskorn lattices was constructed. It is especially a complex manifold, and $G_{\mathbb{Z}}(f_0)$ acts properly discontinuously on it. The quotient $D_{BL}/G_{\mathbb{Z}}$ is a space of isomorphism classes of Brieskorn lattices. There is a natural holomorphic period map

$$BL : M_{\mu}^{mar}(f_0) \rightarrow D_{BL}(f_0).$$

It is an immersion [He6, theorem 12.8] (this refines a weaker result in [SaM]). And it is $G_{\mathbb{Z}}$ -equivariant.

Conjecture 1.2. (a) [He7, conjecture 5.3] *BL is an embedding.*

(b) [He7, conjecture 5.4], [He6, conjecture 12.7], [He1, Kap. 2 d)]) *$LBL : M_{\mu} = M_{\mu}^{mar}/G_{\mathbb{Z}} \rightarrow D_{BL}/G_{\mathbb{Z}}$ is an embedding.*

Part (b) says that the right equivalence class of f is determined by the isomorphism class of $H_0''(f)$. Part (a) would imply part (b). Both are global Torelli type conjectures. Part (b) was proved in [He1] for all simple and unimodal singularities and almost all bimodal singularities, all except three subseries of the eight bimodal series. Therefore for the proof of part (a) in these cases it remains mainly to control $G_{\mathbb{Z}}$ well. But that is surprisingly difficult.

In [He7] the conjectures 1.1 and 1.2 were proved for all simple and those 22 of the 28 exceptional unimodal and bimodal singularities, where all eigenvalues of the monodromy have multiplicity one. In this paper they will be proved for the remaining unimodal and exceptional bimodal singularities, that means, for the simple elliptic and the hyperbolic singularities and for those 6 of the 28 exceptional unimodal and bimodal singularities which had not been treated in [He7].

A priori, logically conjecture 1.1 comes before conjecture 1.2. But the results in [He1] are more concrete and give already some information about the action of $G_{\mathbb{Z}}$ on D_{BL} and M_{μ}^{mar} . Anyway, the main remaining work is a good control of the groups $G_{\mathbb{Z}}$. That presents some unexpected difficulties. For example, we need two surprising generalizations of the number theoretic fact $\mathbb{Z}[e^{2\pi i/a}] \cap S^1 = \{\pm e^{2\pi i k/a} \mid k \in \mathbb{Z}\}$ for $a \in \mathbb{N}$, one is lemma 2.5, the other is related to U_{16} , see remark 4.3.

The groups $G_{\mathbb{Z}}(f_0)$ will be calculated in the sections 3 and 4. Section 2 collects well known background material on the topology of singularities. But it contains also an algebraic lemma 2.5 about automorphisms of monodromy modules. Section 5 collects notions and results from [He7] on marked singularities, the moduli spaces $M_{\mu}^{mar}(f_0)$, the groups $G^{mar}(f_0)$ and the Torelli type conjectures. The sections 6 and 7 give the proofs of the conjectures 1.1 and 1.2 in the cases considered. Section 8 is motivated by the paper [MS] of Milanov and Shen and complements

their results on (transversal) monodromy groups for certain families of simple elliptic singularities. The three principal congruence subgroups $\Gamma(3)$, $\Gamma(4)$ and $\Gamma(6)$ which turn up in [MS] for certain families are shown to turn up also in the biggest possible families.

2. REVIEW ON THE TOPOLOGY OF ISOLATED HYPERSURFACE SINGULARITIES

First, we recall some classical facts and fix some notations. An *isolated hypersurface singularity* (short: *singularity*) is a holomorphic function germ $f : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$ with an isolated singularity at 0. Its *Milnor number*

$$\mu := \dim \mathcal{O}_{\mathbb{C}^{n+1}, 0} / \left(\frac{\partial f}{\partial x_i} \right)$$

is finite. For the following notions and facts compare [AGV2], [Eb2] and (for the notion of an unfolding) [AGV1]. A *good representative* of f has to be defined with some care [Mi][AGV2][Eb2]. It is $f : Y \rightarrow T$ with $Y \subset \mathbb{C}^{n+1}$ a suitable small neighborhood of 0 and $T \subset \mathbb{C}$ a small disk around 0. Then $f : Y' \rightarrow T'$ with $Y' = Y - f^{-1}(0)$ and $T' = T - \{0\}$ is a locally trivial C^∞ -fibration, the *Milnor fibration*. Each fiber has the homotopy type of a bouquet of μ n -spheres [Mi].

Therefore the (reduced for $n = 0$) middle homology groups are $H_n^{(red)}(f^{-1}(\tau), \mathbb{Z}) \cong \mathbb{Z}^\mu$ for $\tau \in T'$. Each comes equipped with an intersection form I , which is a datum of one fiber, a monodromy M_h and a Seifert form L , which come from the Milnor fibration, see [AGV2, I.2.3] for their definitions (for the Seifert form, there are several conventions in the literature, we follow [AGV2]). M_h is a quasiunipotent automorphism, I and L are bilinear forms with values in $\mathbb{Z}$, I is $(-1)^n$ -symmetric, and L is unimodular. L determines M_h and I because of the formulas [AGV2, I.2.3]

$$L(M_h a, b) = (-1)^{n+1} L(b, a), \quad (1)$$

$$I(a, b) = -L(a, b) + (-1)^{n+1} L(b, a). \quad (2)$$

The Milnor lattices $H_n(f^{-1}(\tau), \mathbb{Z})$ for all Milnor fibrations $f : Y' \rightarrow T'$ and then all $\tau \in \mathbb{R}_{>0} \cap T'$ are canonically isomorphic, and the isomorphisms respect M_h , I and L . This follows from Lemma 2.2 in [LR]. These lattices are identified and called *Milnor lattice* $ML(f)$. The group $G_{\mathbb{Z}}$ is

$$G_{\mathbb{Z}} = G_{\mathbb{Z}}(f) := \text{Aut}(ML(f), L) = \text{Aut}(ML(f), M_h, I, L), \quad (3)$$

the second equality is true because L determines M_h and I . We will use the notation $ML(f)_{\mathbb{C}} := ML(f) \otimes_{\mathbb{Z}} \mathbb{C}$, and analogously for other

rings R with $\mathbb{Z} \subset R \subset \mathbb{C}$, and the notations

$$\begin{aligned} Ml(f)_\lambda &:= \ker((M_h - \lambda \text{id})^\mu : Ml(f)_\mathbb{C} \rightarrow Ml(f)_\mathbb{C}) \subset Ml(f)_\mathbb{C}, \\ Ml(f)_{1,\mathbb{Z}} &:= Ml(f)_1 \cap Ml(f) \subset Ml(f), \\ Ml(f)_{\neq 1} &:= \bigoplus_{\lambda \neq 1} Ml(f)_\lambda \subset Ml(f)_\mathbb{C}, \\ Ml(f)_{\neq 1,\mathbb{Z}} &:= Ml(f)_{\neq 1} \cap Ml(f) \subset Ml(f). \end{aligned}$$

The formulas (1) and (2) show $I(a, b) = L((M_h - \text{id})a, b)$. Therefore the eigenspace with eigenvalue 1 of M_h is the radical $\text{Rad}(I) \subset Ml(f)$ of I . By (2) L is $(-1)^{n+1}$ -symmetric on the radical of I .

In the case of a curve singularity ($n = 1$) with r branches, $f = \prod_{j=1}^r f_j$, the radical of I is a $\mathbb{Z}$ -lattice of rank $r - 1$, and it is generated by the classes $l_j \in Ml(f)$ which are obtained by pushing the (correctly oriented) cycles $\partial Y \cap f_j^{-1}(0)$ from the boundary of the fiber $f^{-1}(0)$ to the boundary of the fiber $f^{-1}(\tau)$. Then

$$l_1 + \dots + l_r = 0, \quad (4)$$

$$\begin{aligned} L(l_i, l_j) &= \text{intersection number of } (f_i, f_j) \quad \text{for } i \neq j, \\ \text{so } L(l_i, l_j) &> 0 \quad \text{for } i \neq j, \end{aligned} \quad (5)$$

$$l_1, \dots, \widehat{l_j}, \dots, l_r \quad \text{is a } \mathbb{Z}\text{-basis of } \text{Rad}(I) \text{ for any } j. \quad (6)$$

Kaenders proved the following result using Selling reduction. It will be useful for the calculation of $\text{Aut}(\text{Rad}(I), L)$, because it implies that any automorphism of $(\text{Rad}(I), L)$ maps the set $\{l_1, \dots, l_r\}$ to itself or to minus itself.

Theorem 2.1. [Ka] *In the case of a curve singularity as above, the set $\{l_1, \dots, l_r\}$ is up to a common sign uniquely determined by the properties (4), (5) and (6). So it is up to a common sign determined by the pair $(\text{Rad}(I), L)$. Furthermore, L is negative definite on $\text{Rad}(I)$.*

Examples 2.2. In the following three examples $\underline{l} = (l_1, \dots, l_r)$, and $L(\underline{l}^t, \underline{l}) = (L(l_i, l_j))$. Then $(L(l_i, l_j))_{1 \leq i, j \leq r-1}$ is the matrix of L on $\text{Rad}(I)$ with respect to the $\mathbb{Z}$ -basis $l_1, \dots, l_{r-1}$. The examples (ii) and (iii) will be useful in section 4, the example (i) is an alternative to a calculation in the proof of theorem 8.4 in [He7].

$$(i) \ D_{2m} : f = x^{2m-1} + xy^2 = x(x^{m-1} - iy)(x^{m-1} + iy),$$

$$L(\underline{l}^t, \underline{l}) = \begin{pmatrix} -2 & 1 & 1 \\ 1 & -m & m-1 \\ 1 & m-1 & -m \end{pmatrix}, \quad (L(l_i, l_j))_{1 \leq i, j \leq 2} = \begin{pmatrix} -2 & 1 \\ 1 & -m \end{pmatrix}.$$

Obviously $|\operatorname{Aut}(\operatorname{Rad}(I), L)| = 12$ in the case $m = 2$, and $|\operatorname{Aut}(\operatorname{Rad}(I), L)| = 4$ in the cases $m \geq 3$.

$$(ii) \ Z_{12} : f = x^3y + xy^4 = xy(x^2 + y^3),$$

$$L(\underline{l}^t, \underline{l}) = \begin{pmatrix} -4 & 1 & 3 \\ 1 & -3 & 2 \\ 3 & 2 & -5 \end{pmatrix}, \quad (L(l_i, l_j))_{1 \leq i, j \leq 2} = \begin{pmatrix} -4 & 1 \\ 1 & -3 \end{pmatrix}.$$

Obviously $\operatorname{Aut}(\operatorname{Rad}(I), L) = \{\pm \operatorname{id}\}$.

$$(iii) \ Z_{18} : f = x^3y + xy^6 = xy(x^2 + y^5),$$

$$L(\underline{l}^t, \underline{l}) = \begin{pmatrix} -6 & 1 & 5 \\ 1 & -3 & 2 \\ 5 & 2 & -7 \end{pmatrix}, \quad (L(l_i, l_j))_{1 \leq i, j \leq 2} = \begin{pmatrix} -6 & 1 \\ 1 & -3 \end{pmatrix}.$$

Obviously $\operatorname{Aut}(\operatorname{Rad}(I), L) = \{\pm \operatorname{id}\}$.

Finally, in section 3, distinguished bases will be used. Again, good references for them are [AGV2] and [Eb2]. We sketch their construction and properties.

One can choose a *universal unfolding* of f , a *good representative* F of it with base space $M \subset \mathbb{C}^\mu$, and a generic parameter $t \in M$. Then $F_t : Y_t \rightarrow T$ with $T \subset \mathbb{C}$ the same disk as above and $Y_t \subset \mathbb{C}^{n+1}$ is a *morsification* of f . It has μ A_1 -singularities, and their critical values $u_1, \dots, u_\mu \in T$ are pairwise different. Their numbering is also a choice. Now choose a value $\tau \in T \cap \mathbb{R}_{>0} - \{u_1, \dots, u_\mu\}$ and a *distinguished system of paths*. That is a system of μ paths γ_j , $j = 1, \dots, \mu$, from u_j to τ which do not intersect except at τ and which arrive at τ in clockwise order. Finally, shift from the A_1 singularity above each value u_j the (up to sign unique) vanishing cycle along γ_j to the Milnor fiber $Ml(f) = H_n(f^{-1}(\tau), \mathbb{Z})$, and call the image δ_j .

The tuple $(\delta_1, \dots, \delta_\mu)$ forms a $\mathbb{Z}$ -basis of $Ml(f)$. All such bases are called *distinguished bases*. They form one orbit of an action of a semidirect product $\operatorname{Br}_\mu \ltimes \{\pm 1\}^\mu$. Here Br_μ is the braid group with μ strings, see [AGV2] or [Eb2] for its action. The *sign change group* $\{\pm 1\}^\mu$ acts simply by changing the signs of the entries of the tuples $(\delta_1, \dots, \delta_\mu)$. The members of the distinguished bases are called *vanishing cycles*.

The *Stokes matrix* S of a distinguished basis is defined as the upper triangular matrix in $M(\mu \times \mu, \mathbb{Z})$ with 1's on the diagonal and with

$$S_{ij} := (-1)^{n(n+1)/2} \cdot I(\delta_i, \delta_j) \quad \text{for all } i, j \text{ with } i < j.$$

The *Coxeter-Dynkin diagram* of a distinguished basis encodes S in a geometric way. It has μ vertices which are numbered from 1 to μ . Between two vertices i and j with $i < j$ one draws

no edge if $S_{ij} = 0$,
 $|S_{ij}|$ edges if $S_{ij} < 0$,
 S_{ij} dotted edges if $S_{ij} > 0$.

Coxeter-Dynkin diagrams of many singularities were calculated by A'Campo, Ebeling, Gabrielov and Gusein-Zade. Some of them can be found in [Ga], [Eb1] and [Eb2].

Example 2.3. The hyperbolic singularities of type T_{pqr} with $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$ and the simple elliptic singularities of types $\tilde{E}_6 = T_{333}$, $\tilde{E}_7 = T_{442}$ and $\tilde{E}_8 = T_{632}$ have distinguished bases with the Coxeter-Dynkin diagrams in figure 1 [Ga].




FIGURE 1. A Coxeter-Dynkin diagram of the singularities of type T_{pqr}

The Picard-Lefschetz transformation on $MI(f)$ of a vanishing cycle δ is

$$s_\delta(b) := b - (-1)^{n(n+1)/2} \cdot I(\delta, b) \cdot \delta. \quad (7)$$

The monodromy M_h is

$$M_h = s_{\delta_1} \circ \dots \circ s_{\delta_\mu} \quad (8)$$

for any distinguished basis $\underline{\delta} = (\delta_1, \dots, \delta_\mu)$. The matrices of the monodromy, Seifert form and intersection form with respect to a distinguished basis $\underline{\delta}$ are given by the following formulas.

$$M_h(\underline{\delta}) = \underline{\delta} \cdot (-1)^{n+1} \cdot S^{-1} S^t, \quad (9)$$

$$I(\underline{\delta}^t, \underline{\delta}) = (-1)^{n(n+1)/2} \cdot (S + (-1)^n S^t), \quad (10)$$

$$L(\underline{\delta}^t, \underline{\delta}) = (-1)^{(n+1)(n+2)/2} \cdot S^t. \quad (11)$$

Remark 2.4. The Stokes matrix S of a distinguished basis is related to the matrix V in [Eb2, Korollar 5.3 (i)] by the formula $V = L(\underline{\delta}^t, \underline{\delta}) = (-1)^{(n+1)(n+2)/2} \cdot S^t$. Thus V is *lower* triangular, not *upper* triangular, contrary to the claim in [Eb2, Korollar 5.3 (i)]. The matrix of Var^{-1} for a distinguished basis $\underline{\delta}$ and its dual basis $\underline{\delta}^*$ is $(-1)^{(n+1)(n+2)/2} S$, namely

$$\text{Var}^{-1}(\underline{\delta}) = \underline{\delta}^* \cdot (-1)^{(n+1)(n+2)/2} \cdot S.$$

But then the matrix for the Seifert form L with $L(a, b) = (\text{Var}^{-1}(a))(b)$ is

$$L(\underline{\delta}^t, \underline{\delta}) = \text{Var}^{-1}(\underline{\delta}^t)(\underline{\delta}) = (-1)^{(n+1)(n+2)/2} S^t.$$

In [AGV2, I.2.5] this problem does not arise because there the matrix of a bilinear form with respect to a basis is the transpose of the usual matrix [AGV2, page 45]. This is applied to the matrices of I and L .

A result of Thom and Sebastiani compares the Milnor lattices and monodromies of the singularities $f = f(x_0, \dots, x_n), g = g(y_0, \dots, y_m)$ and $f + g = f(x_0, \dots, x_n) + g(x_{n+1}, \dots, x_{n+m+1})$. There are extensions by Deligne for the Seifert form and by Gabrielov for distinguished bases. All results can be found in [AGV2, I.2.7]. They are restated here. There is a canonical isomorphism

$$\Phi : \text{Ml}(f + g) \xrightarrow{\cong} \text{Ml}(f) \otimes \text{Ml}(g), \quad (12)$$

$$\text{with } M_h(f + g) \cong M_h(f) \otimes M_h(g) \quad (13)$$

$$\text{and } L(f + g) \cong (-1)^{(n+1)(m+1)} \cdot L(f) \otimes L(g). \quad (14)$$

If $\underline{\delta} = (\delta_1, \dots, \delta_{\mu(f)})$ and $\underline{\gamma} = (\gamma_1, \dots, \gamma_{\mu(g)})$ are distinguished bases of f and g with Stokes matrices $S(f)$ and $S(g)$, then

$$\Phi^{-1}(\delta_1 \otimes \gamma_1, \dots, \delta_1 \otimes \gamma_{\mu(g)}, \delta_2 \otimes \gamma_1, \dots, \delta_2 \otimes \gamma_{\mu(g)}, \dots, \delta_{\mu(f)} \otimes \gamma_1, \dots, \delta_{\mu(f)} \otimes \gamma_{\mu(g)})$$

is a distinguished basis of $\text{Ml}(f + g)$, that means, one takes the vanishing cycles $\Phi^{-1}(\delta_i \otimes \gamma_j)$ in the lexicographic order. Then by (11) and (14), the matrix

$$S(f + g) = S(f) \otimes S(g) \quad (15)$$

(where the tensor product is defined so that it fits to the lexicographic order) is the Stokes matrix of this distinguished basis.

In the special case $g = x_{n+1}^2$, the function germ $f + g = f(x_0, \dots, x_n) + x_{n+1}^2 \in \mathcal{O}_{\mathbb{C}^{n+2}, 0}$ is called *stabilization* or *suspension* of f . As there are only two isomorphisms $\text{Ml}(x_{n+1}^2) \rightarrow \mathbb{Z}$, and they differ by a sign, there are two equally canonical isomorphisms $\text{Ml}(f) \rightarrow \text{Ml}(f + x_{n+1}^2)$, and they differ just by a sign. Therefore automorphisms and bilinear forms

on $Gl(f)$ can be identified with automorphisms and bilinear forms on $Gl(f + x_{n+1}^2)$. In this sense

$$L(f + x_{n+1}^2) = (-1)^n \cdot L(f) \quad \text{and} \quad M_h(f + x_{n+1}^2) = -M_h(f) \quad (16)$$

[AGV2, I.2.7], and $G_{\mathbb{Z}}(f + x_{n+1}^2) = G_{\mathbb{Z}}(f)$. The Stokes matrix S does not change under stabilization.

The following algebraic lemma from [He7] will be very useful in the sections 3 and 4. It can be seen as a generalization of the number theoretic fact $\mathbb{Z}[e^{2\pi i/a}] \cap S^1 = \{\pm e^{2\pi i k/a} \mid k \in \mathbb{Z}\}$.

Lemma 2.5. [He7, lemma 8.2] *Let H be a free $\mathbb{Z}$ -module of finite rank μ , and $H_{\mathbb{C}} := H \otimes_{\mathbb{Z}} \mathbb{C}$. Let $M_h : H \rightarrow H$ be an automorphism of finite order, called monodromy, with three properties:*

- (i) *Each eigenvalue has multiplicity 1.*
Denote $H_{\lambda} := \ker(M_h - \lambda \cdot \text{id} : H_{\mathbb{C}} \rightarrow H_{\mathbb{C}})$.
- (ii) *Denote $\text{Ord} := \{\text{ord } \lambda \mid \lambda \text{ eigenvalue of } M_h\} \subset \mathbb{Z}_{\geq 1}$. There exist four sequences $(m_i)_{i=1, \dots, |\text{Ord}|}$, $(j(i))_{i=2, \dots, |\text{Ord}|}$, $(p_i)_{i=2, \dots, |\text{Ord}|}$, $(k_i)_{i=2, \dots, |\text{Ord}|}$ of numbers in $\mathbb{Z}_{\geq 1}$ and two numbers $i_1, i_2 \in \mathbb{Z}_{\geq 1}$ with $i_1 \leq i_2 \leq |\text{Ord}|$ and with the properties:*
 $\text{Ord} = \{m_1, \dots, m_{|\text{Ord}|}\},$
 p_i *is a prime number, $p_i = 2$ for $i_1 + 1 \leq i \leq i_2$, $p_i \geq 3$ else,*
 $j(i) = i - 1$ *for $i_1 + 1 \leq i \leq i_2$, $j(i) < i$ else,*
 $m_i = m_{j(i)} / p_i^{k_i}.$
- (iii) *A cyclic generator $a_1 \in H$ exists, that means,*

$$H = \bigoplus_{i=0}^{\mu-1} \mathbb{Z} \cdot M_h^i(a_1).$$

Finally, let I be an M_h -invariant nondegenerate bilinear form (not necessarily (± 1) -symmetric) on $\bigoplus_{\lambda \neq \pm 1} H_{\lambda}$ with values in $\mathbb{C}$. Then

$$\text{Aut}(H, M_h, I) = \{\pm M_h^k \mid k \in \mathbb{Z}\}.$$

3. THE GROUP $G_{\mathbb{Z}}$ FOR THE SIMPLE ELLIPTIC AND THE HYPERBOLIC SINGULARITIES

The simple elliptic and the hyperbolic singularities are 1-parameter families of singularities, which had been classified by Arnold [AGV1]. For each triple $(p, q, r) \in \mathbb{N}_{\geq 2}^3$ with $p \geq q \geq r$ and $\kappa := \frac{1}{p} + \frac{1}{q} + \frac{1}{r} \leq 1$ one has one family, denoted T_{pqr} . The hyperbolic singularities are those with $\kappa < 1$, the simple elliptic are those with $\kappa = 1$. For the three families of simple elliptic singularities also other symbols are used, $T_{333} = \tilde{E}_6$, $T_{442} = \tilde{E}_7$, $T_{632} = \tilde{E}_8$. The singularities of types T_{pqr} with

$r = 2$ exist as curve singularities, all others as surface singularities. Normal forms will be discussed in section 6. Here for each family the group $G_{\mathbb{Z}} = \text{Aut}(Ml(f), L)$ will be analyzed. The result in Theorem 3.1 is completely explicit in the case $\kappa < 1$ and partly explicit in the case $\kappa = 1$. The whole section is devoted to its proof. Besides

$$\kappa := \frac{1}{p} + \frac{1}{q} + \frac{1}{r}, \quad \text{also} \quad \chi := \text{lcm}(p, q, r) \in \mathbb{N}$$

will be used. A singularity of type T_{pqr} has Milnor number $\mu = p + q + r - 1$, and its monodromy has the characteristic polynomial

$$\frac{t^p - 1}{t - 1} \cdot \frac{t^q - 1}{t - 1} \cdot \frac{t^r - 1}{t - 1} \cdot (t - 1)^2.$$

Theorem 3.1. *Consider a surface singularity f of type T_{pqr} (with $\kappa \leq 1$) with Milnor lattice $Ml(f)$, monodromy M_h , intersection form I and Seifert form L .*

(a) *Then $\dim Ml(f)_1 = 2$, $\text{rank Rad}(I) = 1$ if $\kappa < 1$ and $= 2$ if $\kappa = 1$. Choose a $\mathbb{Z}$ -basis b_1, b_2 of $Ml(f)_{1, \mathbb{Z}}$ with $b_1 \in \text{Rad}(I)$ and $L(b_1, b_2) \leq 0$. Then*

$$L(b_1, b_2) = -\chi \quad \text{and} \quad M_h b_2 = b_2 + \chi(\kappa - 1) \cdot b_1. \quad (17)$$

(b) *The restriction map $G_{\mathbb{Z}} \rightarrow \text{Aut}(Ml(f)_{1, \mathbb{Z}}, L)$ is surjective. Denote by $T \in \text{Aut}(Ml(f)_{1, \mathbb{Z}})$ the automorphism with $T(b_1) = b_1$ and $T(b_2) = b_2 + b_1$. Denote $\underline{b} := (b_1, b_2)$.*

$$\begin{aligned} \text{Aut}(Ml(f)_{1, \mathbb{Z}}, L) &= \{ \underline{b} \mapsto \underline{b} \cdot A \mid A \in SL(2, \mathbb{Z}) \} \\ &\cong SL(2, \mathbb{Z}) \quad \text{if} \quad \kappa = 1, \end{aligned} \quad (18)$$

$$\text{Aut}(Ml(f)_{1, \mathbb{Z}}, L) = \{ \pm T^k \mid k \in \mathbb{Z} \} \quad \text{if} \quad \kappa < 1. \quad (19)$$

(c) *The group $G_{\mathbb{Z}}$ for $\kappa < 1$ and the subgroup $\{g \in G_{\mathbb{Z}} \mid g(b_1) = \pm b_1\} \subset G_{\mathbb{Z}}$ for $\kappa = 1$ will be described explicitly, except for the part U_2 , see below. There is a monodromy invariant decomposition*

$$Ml(f)_{\neq 1} = Ml_{\mathbb{C}}^{(1)} \oplus Ml_{\mathbb{C}}^{(2)} \oplus Ml_{\mathbb{C}}^{(3)} \quad (20)$$

such that the characteristic polynomial of $M_h|_{Ml^{(j)}}$ is

$$\frac{t^p - 1}{t - 1}, \quad \frac{t^q - 1}{t - 1}, \quad \frac{t^r - 1}{t - 1} \quad \text{for } j = 1, 2, 3 \quad (21)$$

and such that the following holds.

$$\left\{ g \in G_{\mathbb{Z}} \mid g(b_1) = \pm b_1 \right\} = \left. \begin{array}{ll} G_{\mathbb{Z}} & \text{for } \kappa < 1 \\ & \text{for } \kappa = 1 \end{array} \right\} = (U_1 \rtimes U_2) \times \{\pm \text{id}\}, \quad (22)$$

where U_1 is the infinite subgroup of $G_{\mathbb{Z}}$

$$U_1 = \{T^\delta \times (M_h|_{M_{\mathbb{C}}^{(1)}})^\alpha \times (M_h|_{M_{\mathbb{C}}^{(2)}})^\beta \times (M_h|_{M_{\mathbb{C}}^{(3)}})^\gamma \mid (\delta, \alpha, \beta, \gamma) \in \mathbb{Z} \times \mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r \text{ with } \frac{\alpha}{p} + \frac{\beta}{q} + \frac{\gamma}{r} \equiv \frac{\delta}{\chi} \pmod{1}\} \quad (23)$$

and where U_2 is a finite subgroup of $G_{\mathbb{Z}}$ with

$$U_2 \begin{cases} = \{\text{id}\} & \text{if } p > q > r, \\ \cong S_2 & \text{if } p = q > r \text{ or } p > q = r, \\ \cong S_3 & \text{if } p = q = r. \end{cases} \quad (24)$$

which consists of certain automorphisms which act trivially on $Ml(f)_1$ and which permute those of the subspaces $Ml_{\mathbb{C}}^{(j)}$ which have equal dimension.

Proof: Choose a distinguished basis $\underline{\delta} = (\delta_1, \dots, \delta_\mu)$ with the Coxeter-Dynkin diagram in example 2.3. Then the monodromy matrix M_M with $M_h(\underline{\delta}) = \underline{\delta} \cdot M_M$ can be calculated either with (8) or with (9). It had been calculated in [He1] with (8), and it is (here all not specified entries are 0)

$$M_M = \begin{pmatrix} M_1 & & & M_8 \\ & M_2 & & M_9 \\ & & M_3 & M_{10} \\ M_5 & M_6 & M_7 & M_4 \end{pmatrix} \quad (25)$$

with the following blocks,

$$M_1 = \begin{pmatrix} 0 & & -1 \\ 1 & \ddots & -1 \\ & \ddots & 0 & -1 \\ & & 1 & -1 \end{pmatrix} \in M((p-1) \times (p-1), \mathbb{Z}),$$

$$M_2 \in M((q-1) \times (q-1), \mathbb{Z}) \text{ and}$$

$$M_3 \in M((r-1) \times (r-1), \mathbb{Z}) \text{ are defined analogously,}$$

$$M_4 = \begin{pmatrix} 3 & 2 \\ -2 & -1 \end{pmatrix},$$

and $M_5, M_6, M_7, M_8, M_9, M_{10}$ are of suitable sizes with all entries except the following being 0,

$$(M_5)_{11} = (M_6)_{11} = (M_7)_{11} = -1, \quad (M_5)_{21} = (M_6)_{21} = (M_7)_{21} = 1,$$

$$(M_8)_{11} = (M_8)_{12} = (M_9)_{11} = (M_9)_{12} = (M_{10})_{11} = (M_{10})_{12} = 1.$$

Define

$$\tilde{b}_1 := \delta_{\mu-1} - \delta_\mu, \quad (26)$$

$$\begin{aligned} \tilde{b}_2 := \chi \cdot & \left(\sum_{i=1}^{p-1} \frac{p-i}{p} \delta_i + \sum_{i=1}^{q-1} \frac{q-i}{q} \delta_{p-1+i} \right. \\ & \left. + \sum_{i=1}^{r-1} \frac{r-i}{r} \delta_{p+q-2+i} + \delta_{\mu-1} \right). \end{aligned} \quad (27)$$

Then one calculates

$$M_h(\tilde{b}_1) = \tilde{b}_1, \quad M_h(\tilde{b}_2) = \tilde{b}_2 + \chi(\kappa - 1) \cdot \tilde{b}_1, \quad (28)$$

and with (11), which is here $L(\underline{\delta}^t, \underline{\delta}) = S^t$,

$$\begin{pmatrix} L(\tilde{b}_1, \tilde{b}_1) & L(\tilde{b}_1, \tilde{b}_2) \\ L(\tilde{b}_2, \tilde{b}_1) & L(\tilde{b}_2, \tilde{b}_2) \end{pmatrix} = \begin{pmatrix} 0 & -\chi \\ \chi & \frac{\chi^2}{2}(\kappa - 1) \end{pmatrix}. \quad (29)$$

By (28), $\tilde{b}_1, \tilde{b}_2$ is a $\mathbb{Q}$ -basis of $Ml(f)_{1,\mathbb{Q}}$ and M_h is on $Ml(f)_1$ semisimple if $\kappa = 1$ and it has a 2×2 Jordan block if $\kappa < 1$ (of course, this is well known). From the coefficients one sees that $\tilde{b}_1, \tilde{b}_2$ is a $\mathbb{Z}$ -basis of $Ml(f)_{1,\mathbb{Z}}$. Here it is important that the coefficients of $\tilde{b}_2$ have greatest common divisor 1. As the equations (17) hold for $\tilde{b}_1, \tilde{b}_2$, they hold for any basis b_1, b_2 as in (a).

(b) If $\kappa = 1$ then (29) shows that L is on $Ml(f)_{1,\mathbb{Z}}$ up to the factor χ the standard symplectic form. Therefore (18) holds. If $\kappa < 1$ then (29) shows (19).

The restriction map $G_{\mathbb{Z}} \rightarrow \text{Aut}(Ml(f)_{1,\mathbb{Z}})$ contains T . This follows from (22) (whose proof below does not use this fact), because obviously $(\delta, \alpha, \beta, \gamma)$ as in (23) with $\delta = 1$ exist.

This shows (b) in the case $\kappa < 1$. In the case $\kappa = 1$ we did not calculate lifts to $G_{\mathbb{Z}}$ of other elements of $\text{Aut}(Ml(f)_{1,\mathbb{Z}}, L)$. In this case the surjectivity of $G_{\mathbb{Z}} \rightarrow \text{Aut}(Ml(f)_{1,\mathbb{Z}}, L)$ follows in two ways: It follows from [Kl, III.2.6], and it follows from calculations in [He1], which are discussed in the proof of theorem 6.1.

(c) We will prove (c) for the special choice $\tilde{b}_1, \tilde{b}_2$. Then (c) holds for any b_1, b_2 as in (a) because by the surjectivity of the map $G_{\mathbb{Z}} \rightarrow \text{Aut}(Ml(f)_{1,\mathbb{Z}}, L)$, an element $g \in G_{\mathbb{Z}}$ with $g(\tilde{b}_1) = b_1, g(\tilde{b}_2) = b_2$ exists.

Define

$$Ml_{\mathbb{Z}}^{[1]} := \mathbb{Z}\tilde{b}_1 \oplus \bigoplus_{i=1}^{p-1} \mathbb{Z}\delta_i, \quad Ml_{\mathbb{C}}^{[1]} := Ml_{\mathbb{Z}}^{[1]} \otimes_{\mathbb{Z}} \mathbb{C}, \quad (30)$$

$$Ml_{\mathbb{C}}^{(1)} := Ml_{\mathbb{C}}^{[1]} \cap Ml(f)_{\neq 1}, \quad (31)$$

and analogously $Ml_{\mathbb{Z}}^{[2]}, Ml_{\mathbb{C}}^{[2]}, Ml_{\mathbb{C}}^{(2)}$ and $Ml_{\mathbb{Z}}^{[3]}, Ml_{\mathbb{C}}^{[3]}, Ml_{\mathbb{C}}^{(3)}$.

A look at the matrix M_M shows the following.

$$M_h : \delta_1 + \tilde{b}_1 \mapsto \delta_2 \mapsto \dots \mapsto \delta_{p-1} \mapsto -(\delta_1 + \dots + \delta_{p-1}) \mapsto \delta_1 + \tilde{b}_1. \quad (32)$$

Therefore $Ml_{\mathbb{Z}}^{[1]}$ is a cyclic M_h -module with characteristic polynomial $t^p - 1$, and $Ml_{\mathbb{C}}^{[1]} = \mathbb{C}\tilde{b}_1 \oplus Ml_{\mathbb{C}}^{(1)}$, and M_h on $Ml_{\mathbb{C}}^{(1)}$ has the characteristic polynomial $(t^p - 1)/(t - 1)$.

Lemma 2.5 applies and shows

$$\text{Aut}(Ml_{\mathbb{Z}}^{[1]}, L) = \{\pm(M_h|_{Ml_{\mathbb{Z}}^{[1]}})^{\alpha} \mid \alpha \in \{0, 1, \dots, p-1\}\}. \quad (33)$$

Finally, M_h, I and L are well defined on the quotient lattice $Ml_{\mathbb{Z}}^{[1]}/\mathbb{Z}\tilde{b}_1$, and $(Ml_{\mathbb{Z}}^{[1]}/\mathbb{Z}\tilde{b}_1, -I)$ is a root lattice of type A_{p-1} . The last statement follows immediately from the part of the Coxeter-Dynkin diagram which corresponds to $\delta_1, \dots, \delta_{p-1}$.

$Ml_{\mathbb{Z}}^{[2]}$ and $Ml_{\mathbb{Z}}^{[3]}$ have the same properties as $Ml_{\mathbb{Z}}^{[1]}$, with q respectively r instead of p .

Now

$$Ml(f)_{\neq 1} = Ml_{\mathbb{C}}^{(1)} \oplus Ml_{\mathbb{C}}^{(2)} \oplus Ml_{\mathbb{C}}^{(3)}$$

is clear. The $\mathbb{Z}$ -lattice

$$Ml_{\mathbb{Z}}^{[1]} + Ml_{\mathbb{Z}}^{[2]} + Ml_{\mathbb{Z}}^{[3]} = \mathbb{Z}\tilde{b}_1 \oplus \bigoplus_{i=1}^{\mu-2} \mathbb{Z}\delta_i = (\mathbb{C}\tilde{b}_1 \oplus Ml(f)_{\neq 1}) \cap Ml(f)$$

is a primitive sublattice of $Ml(f)$ of rank $\mu - 1$. Any $g \in G_{\mathbb{Z}}$ with $g(\tilde{b}_1) = \pm\tilde{b}_1$ maps it to itself, because it maps $\mathbb{C}\tilde{b}_1$ and $Ml(f)_{\neq 1}$ and $Ml(f)$ to themselves. g maps also the quotient lattice

$$(Ml_{\mathbb{Z}}^{[1]} + Ml_{\mathbb{Z}}^{[2]} + Ml_{\mathbb{Z}}^{[3]})/\mathbb{Z}\tilde{b}_1 = Ml_{\mathbb{Z}}^{[1]}/\mathbb{Z}\tilde{b}_1 \oplus Ml_{\mathbb{Z}}^{[2]}/\mathbb{Z}\tilde{b}_1 \oplus Ml_{\mathbb{Z}}^{[3]}/\mathbb{Z}\tilde{b}_1$$

to itself. But this is together with $-I$ an orthogonal sum of lattices of types A_{p-1} , A_{q-1} and A_{r-1} . Therefore g can only permute the summands, and only those summands of equal rank.

If $p = q$, a special element $\sigma_{12} \in G_{\mathbb{Z}}$ is given by

$$\begin{aligned} \sigma_{12}(\delta_i) &= \delta_{p-1+i}, & \sigma_{12}(\delta_{p-1+i}) &= \delta_i \quad \text{for } 1 \leq i \leq p-1, \\ \sigma_{12}(\delta_j) &= \delta_j \quad \text{for } p+q-2 \leq j \leq \mu. \end{aligned}$$

$\sigma_{12} \in G_{\mathbb{Z}}$ follows immediately from the symmetry of the Coxeter-Dynkin diagram. Similarly $\sigma_{23} \in G_{\mathbb{Z}}$ is defined if $q = r$. In any case, these elements generate a subgroup $U_2 \subset G_{\mathbb{Z}}$ with the properties in (c).

Therefore, starting with an arbitrary element $\tilde{g} \in G_{\mathbb{Z}}$ if $\kappa < 1$ respectively $\tilde{g} \in \{g \in G_{\mathbb{Z}} \mid g(\tilde{b}_1) = \pm \tilde{b}_1\}$ if $\kappa = 1$, one can compose it with $\pm \text{id}$ and an element of U_2 , and one obtains an element $g \in G_{\mathbb{Z}}$ with $g(\tilde{b}_1) = \tilde{b}_1$ and $g(Ml_{\mathbb{Z}}^{[j]}) = Ml_{\mathbb{Z}}^{[j]}$ for $j = 1, 2, 3$. Then $g|_{Ml_{\mathbb{Z}}^{[1]}} = (M_h|_{Ml_{\mathbb{Z}}^{[1]}})^{\alpha}$ for a unique $\alpha \in \{0, 1, \dots, p-1\}$, and similarly with $\beta \in \{0, 1, \dots, q-1\}$ and $\gamma \in \{0, 1, \dots, r-1\}$ for $Ml_{\mathbb{Z}}^{[2]}$ and $Ml_{\mathbb{Z}}^{[3]}$. Also $g(\tilde{b}_2) = \tilde{b}_2 + \delta \tilde{b}_1$ for some $\delta \in \mathbb{Z}$. One calculates, observing (32),

$$M_h \left(\sum_{i=1}^{p-1} \frac{p-i}{p} \delta_i \right) = \left(\sum_{i=1}^{p-1} \frac{p-i}{p} \delta_i \right) - (\delta_1 + \tilde{b}_1) + \frac{1}{p} \tilde{b}_1, \quad (34)$$

$$M_h^{\alpha} \left(\sum_{i=1}^{p-1} \frac{p-i}{p} \delta_i \right) = \left(\sum_{i=1}^{p-1} \frac{p-i}{p} \delta_i \right) - \left(\tilde{b}_1 + \sum_{i=1}^{\alpha} \delta_i \right) + \frac{\alpha}{p} \tilde{b}_1. \quad (35)$$

The definition (27) of $\tilde{b}_2$ shows

$$-\delta_{\mu-1} = -\frac{1}{\chi} \tilde{b}_2 + \sum_{i=1}^{p-1} \frac{p-i}{p} \delta_i + \sum_{i=1}^{q-1} \frac{q-i}{q} \delta_{p-1+i} + \sum_{i=1}^{r-1} \frac{r-i}{r} \delta_{p+q-2+i}, \quad (36)$$

(35) gives

$$\begin{aligned} g(-\delta_{\mu-1}) &= -\delta_{\mu-1} + \left(\frac{-\delta}{\chi} + \frac{\alpha}{p} + \frac{\beta}{q} + \frac{\gamma}{r} \right) \cdot \tilde{b}_1 \\ &\quad - \left(\tilde{b}_1 + \sum_{i=1}^{\alpha} \delta_i \right) - \left(\tilde{b}_1 + \sum_{i=p}^{p-1+\beta} \delta_i \right) - \left(\tilde{b}_1 + \sum_{i=p+q-1}^{p+q-2+\gamma} \delta_i \right). \end{aligned} \quad (37)$$

Therefore

$$\begin{aligned} \frac{\alpha}{p} + \frac{\beta}{q} + \frac{\gamma}{r} &\equiv \frac{\delta}{\chi} \pmod{1} \\ \text{and } g &= T^{\delta} \times (M_h|_{Ml_{\mathbb{C}}^{(1)}})^{\alpha} \times (M_h|_{Ml_{\mathbb{C}}^{(2)}})^{\beta} \times (M_h|_{Ml_{\mathbb{C}}^{(3)}})^{\gamma}. \end{aligned} \quad (38)$$

Thus $g \in U_1$, so $G_{\mathbb{Z}} \subset (U_1 \rtimes U_2) \times \{\pm \text{id}\}$.

Vice versa, we have to show $U_1 \subset G_{\mathbb{Z}}$. Fix a $g \in U_1$. It respects the decomposition

$$Ml(f)_{\mathbb{C}} = Ml(f)_1 \oplus Ml_{\mathbb{C}}^{(1)} \oplus Ml_{\mathbb{C}}^{(2)} \oplus Ml_{\mathbb{C}}^{(3)}.$$

This is a left and right orthogonal decomposition with respect to the Seifert form L . The restriction of g to each of the four blocks respects

L there, so $g \in \text{Aut}(Ml(f)_{\mathbb{C}}, L)$. It restricts on $Ml_{\mathbb{C}}^{[1]}$ to M_h^α , so it maps the lattice $Ml_{\mathbb{Z}}^{[1]}$ to itself, and analogously the lattices $Ml_{\mathbb{Z}}^{[2]}$ and $Ml_{\mathbb{Z}}^{[3]}$, thus also the sum $Ml_{\mathbb{Z}}^{[1]} + Ml_{\mathbb{Z}}^{[2]} + Ml_{\mathbb{Z}}^{[3]}$. This sum is a primitive sublattice of $Ml(f)$ of rank $\mu - 1$ with

$$Ml(f) = \left(Ml_{\mathbb{Z}}^{[1]} + Ml_{\mathbb{Z}}^{[2]} + Ml_{\mathbb{Z}}^{[3]} \right) \oplus \mathbb{Z}\delta_{\mu-1}.$$

The calculation above of $g(-\delta_{\mu-1})$ shows $g(\delta_{\mu-1}) \in Ml(f)$ and $g(\delta_{\mu-1}) \equiv \delta_{\mu-1}$ modulo the sublattice. Therefore $g \in G_{\mathbb{Z}}$. $\square$

4. THE GROUP $G_{\mathbb{Z}}$ FOR 6 OF THE 28 EXCEPTIONAL UNIMODAL AND BIMODAL SINGULARITIES

The 14 1-parameter families of exceptional unimodal singularities and the 14 2-parameter families of exceptional bimodal singularities had been classified by Arnold. Normal forms can be found in [AGV1]. In [He7, theorem 8.3] the group $G_{\mathbb{Z}}$ was calculated for 22 of the 28 families, namely for those families where all eigenvalues of the monodromy have multiplicity 1. In these cases it turned out that $G_{\mathbb{Z}}$ is simply $\{\pm M_h^k \mid k \in \mathbb{Z}\}$. The proof used lemma 2.5 and that the Milnor lattice is in these cases a cyclic monodromy module.

In this section $G_{\mathbb{Z}}$ will be determined for the remaining 6 of the 28 families. These are the families $Z_{12}, Q_{12}, U_{12}, Z_{18}, Q_{16}, U_{16}$. In these cases, some eigenvalues have multiplicity 2. This is similar to the case of the singularity D_{2m} , which had also been treated in [He7, theorem 8.4]. Also the proof will be similar. It will again use lemma 2.5 and combine that with an additional analysis of the action of $G_{\mathbb{Z}}$ on the sum of the eigenspaces with dimension = 2, see lemma 4.2 below.

This lemma presents a surprise, it points at a funny generalization of the number theoretic fact $\mathbb{Z}[e^{2\pi i/n}] \cap S_1 = \{\pm e^{2\pi i k/n} \mid k \in \mathbb{Z}\}$. Also, the lemma uses at the end a calculation of $G^{mar} \subset G_{\mathbb{Z}}$, which will only come in section 7. The whole section 4 is devoted to the proof of the following theorem.

Theorem 4.1. *In the case of the 6 families of exceptional unimodal and bimodal singularities $Z_{12}, Q_{12}, U_{12}, Z_{18}, Q_{16}$ and U_{16} , the group $G_{\mathbb{Z}}$ is $G_{\mathbb{Z}} = \{\pm M_h^k \mid k \in \mathbb{Z}\} \times U$ with*

$$U \cong \begin{array}{c|c|c|c|c|c|c} & Z_{12} & Q_{12} & U_{12} & Z_{18} & Q_{16} & U_{16} \\ \hline & \{\text{id}\} & S_2 & S_3 & \{\text{id}\} & S_2 & S_3 \end{array} \quad (39)$$

(This is independent of the number of variables, i.e. it does not change under stabilization.)

Proof: Here we consider all 6 families as surface singularities. Their characteristic polynomials p_{ch} have all the property $p_{ch} = p_1 \cdot p_2$ with $p_2|p_1$ and p_1 having only simple roots. They are as follows. Here Φ_m is the cyclotomic polynomial of primitive m -th unit roots.

	Z_{12}	Q_{12}	U_{12}	Z_{18}	Q_{16}	U_{16}
p_{ch}	$\Phi_{22}\Phi_2^2$	$\Phi_{15}\Phi_3^2$	$\Phi_{12}\Phi_6\Phi_4\Phi_2^2$	$\Phi_{34}\Phi_2^2$	$\Phi_{21}\Phi_3^2$	$\Phi_{15}\Phi_5^2$
p_1	$\Phi_{22}\Phi_2$	$\Phi_{15}\Phi_3$	$\Phi_{12}\Phi_6\Phi_4\Phi_2$	$\Phi_{34}\Phi_2$	$\Phi_{21}\Phi_3$	$\Phi_{15}\Phi_5$
p_2	Φ_2	Φ_3	$\Phi_4\Phi_2$	Φ_2	Φ_3	Φ_5

(40)

Orlik [Or] had conjectured that the Milnor lattice of any quasi-homogeneous singularity is a sum of cyclic monodromy modules and that the characteristic polynomials of M_h on the cyclic pieces are $p_1, \dots, p_r$ where $p_{ch} = p_1 \cdot \dots \cdot p_r$ and $p_r|p_{r-1}|\dots|p_1$ and p_1 has simple roots (r and $p_1, \dots, p_r$ are uniquely determined by this). In the case of curve singularities, Michel and Weber [MW] have a proof of this conjecture. In [He1, 3.1] the conjecture was proved (using Coxeter-Dynkin diagrams) for all those quasihomogeneous surface singularities of modality ≤ 2 which are not stabilizations of curve singularities. So especially, the conjecture is true for the families of singularities $Z_{12}, Q_{12}, U_{12}, Z_{18}, Q_{16}, U_{16}$. There are $a_1, a_2 \in \text{Ml}(f)$ with

$$\text{Ml}(f) = \left(\bigoplus_{i=0}^{\deg p_1 - 1} \mathbb{Z} \cdot M_h^i(a_1) \right) \oplus \left(\bigoplus_{i=0}^{\deg p_2 - 1} \mathbb{Z} \cdot M_h^i(a_2) \right) =: B_1 \oplus B_2. \quad (41)$$

Denote

$$B_3 := \ker(p_2(M_h) : \text{Ml}(f)_{\mathbb{C}} \rightarrow \text{Ml}(f)_{\mathbb{C}}) \cap \text{Ml}(f). \quad (42)$$

It is a primitive sublattice of $\text{Ml}(f)$ of rank $2 \deg p_2$. Also,

$$(B_1)_{\mathbb{C}} = \ker((p_1/p_2)(M_h)) \oplus (B_1 \cap B_3)_{\mathbb{C}}, \quad B_2 \subset B_3, \quad (43)$$

and $B_1 \cap B_3$ and B_2 are both M_h -invariant primitive sublattices of B_3 of rank $\deg p_2$. Together they generate B_3 .

Any $g \in G_{\mathbb{Z}}$ with $g|_{B_3} = \pm(M_h|_{B_3})^k$ for some $k \in \mathbb{Z}$ restricts because of (43) to an automorphism of B_1 . Lemma 2.5 applies and shows $g|_{B_1} = \pm(M_h|_{B_1})^l$ for some $l \in \mathbb{Z}$. Now $g|_{B_3} = \pm(M_h|_{B_3})^k$ enforces $k \equiv l \pmod{\text{lcm}(m|\Phi_m|p_2)}$ and $g = \pm M_h^l$. Therefore

$$\{g \in G_{\mathbb{Z}} \mid g|_{B_3} = \pm(M_h|_{B_3})^k \text{ for some } k\} = \{\pm M_h^k \mid k \in \mathbb{Z}\}. \quad (44)$$

Lemma 4.2 (c) determines $\text{Aut}(B_3, L)$,

$$\text{Aut}(B_3, L) = \{\pm(M_h|_{B_3})^k \mid k \in \mathbb{Z}\} \times U \quad \text{with } U \text{ as in (39)}. \quad (45)$$

Lemma 4.2 (d) shows that the map $G_{\mathbb{Z}} \rightarrow \text{Aut}(B_3, L)$ is surjective. Together with (44) this gives (39). $\square$

Lemma 4.2. (a) Let $V_{\mathbb{Z}}$ be a $\mathbb{Z}$ -lattice of rank 2 with a $\mathbb{Z}$ -basis $\underline{b} = (b_1, b_2)$ and a symmetric pairing $L_{\mathbb{Z}}$ given by

$$L_{\mathbb{Z}}(\underline{b}^t, \underline{b}) = \begin{pmatrix} 2 & -1 \\ -1 & m \end{pmatrix} \quad \text{for some } m \in \mathbb{N}_{\geq 2}.$$

Define $\xi := e^{2\pi i/l}$ for some $l \in \{3, 4, 5\}$, where we exclude the cases $(m \geq 3, l = 5)$. Define $V_{\mathbb{C}} := V_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{C}$, $V_{\mathbb{Z}[\xi]} := V_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Z}[\xi] \subset V_{\mathbb{C}}$, and extend $L_{\mathbb{Z}}$ sesquilinearly (=linear $\times$ semilinear) to $V_{\mathbb{C}}$. Then

$$\begin{aligned} & \{r \in V_{\mathbb{Z}[\xi]} \mid L_{\mathbb{C}}(r, r) = 2\} \\ &= \{\pm \xi^k \mid k \in \mathbb{Z}\} \times \{r \in V_{\mathbb{Z}} \mid L_{\mathbb{Z}}(r, r) = 2\}. \end{aligned} \quad (46)$$

In the case $m \geq 3$

$$\begin{aligned} & \{r \in V_{\mathbb{Z}[\xi]} \mid L_{\mathbb{C}}(r, r) = m, r \notin \mathbb{Z}[\xi]b_1\} \\ &= \{\pm \xi^k \mid k \in \mathbb{Z}\} \times \{r \in V_{\mathbb{Z}} \mid L_{\mathbb{Z}}(r, r) = m, r \notin \mathbb{Z}b_1\}. \end{aligned} \quad (47)$$

(b) In the situation of (a)

$$\begin{aligned} \text{Aut}(V_{\mathbb{Z}[\xi]}, L_{\mathbb{C}}) &= \{\pm \xi^k \mid k \in \mathbb{Z}\} \cdot \text{Aut}(V_{\mathbb{Z}}, L_{\mathbb{Z}}) \\ & \quad (\pm \text{id} \text{ lives on both sides, therefore not } \times) \end{aligned} \quad (48)$$

$$\begin{aligned} \text{Aut}(V_{\mathbb{Z}}, L_{\mathbb{Z}}) &\cong \left\{ \pm \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \pm \begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix} \right\} \\ &\cong \{\pm \text{id}\} \times S_2 \quad \text{in the cases } m \geq 3, \end{aligned} \quad (49)$$

$$\begin{aligned} \text{Aut}(V_{\mathbb{Z}}, L_{\mathbb{Z}}) &\cong \text{Aut}(\text{root lattice of type } A_2) \\ &\cong \{\pm \text{id}\} \times S_3 \quad \text{in the case } m = 2. \end{aligned} \quad (50)$$

(c) In the situation of the proof of theorem 4.1

$$\text{Aut}(B_3, L) = \{\pm (M_h|_{B_3})^k \mid k \in \mathbb{Z}\} \times U \quad \text{with } U \text{ as in (39)}. \quad (51)$$

(d) In the situation of the proof of theorem 4.1 the map $G_{\mathbb{Z}} \rightarrow \text{Aut}(B_3, L)$ is surjective.

Proof: (a) An element $r = r_1 b_1 + r_2 b_2$ with $r_1, r_2 \in \mathbb{Z}[\xi]$ satisfies

$$\begin{aligned} L_{\mathbb{C}}(r, r) &= 2|r_1|^2 - (r_1 \bar{r}_2 + \bar{r}_1 r_2) + m|r_2|^2 \\ &= |r_1|^2 + |r_1 - r_2|^2 + (m-1)|r_2|^2. \end{aligned} \quad (52)$$

First consider the cases $l \in \{3, 4\}$. Then $\mathbb{Z}[\xi] \cap \mathbb{R} = \mathbb{Z}$. Then the three absolute values in (52) are non-negative integers. Their sum is 2 if and only if

$$|r_1| = 1, r_2 = 0 \quad \text{in the cases } m \geq 3, \quad (53)$$

$$\left. \begin{aligned} & (|r_1|, |r_2|) \in \{(1, 0), (0, 1), (1, 1)\} \\ & \text{and in the last case } r_1 = r_2 \end{aligned} \right\} \quad \text{in the case } m = 2. \quad (54)$$

In the case $m \geq 3$ and in the case of an $r \notin \mathbb{Z}[\xi]b_1$, the sum of the three absolute values in (53) is m if and only if

$$(r_1 = 0, |r_2| = 1) \text{ or } (r_1 = r_2, |r_1| = 1). \quad (55)$$

Together with $\mathbb{Z}[\xi] \cap S^1 = \{\pm \xi^k \mid k \in \mathbb{Z}\}$ this shows part (a) in the cases $l \in \{3, 4\}$.

It rests to consider the case $(m, l) = (2, 5)$. In that case write

$$r_1 = r_{10} + r_{11}\xi + r_{12}\xi^2 + r_{13}\xi^3, \quad r_2 = r_{20} + r_{21}\xi + r_{22}\xi^2 + r_{23}\xi^3$$

with $r_{ij} \in \mathbb{Z}$. Then

$$\begin{aligned} L_{\mathbb{C}}(r, r) &= 2|r_1|^2 + 2|r_2|^2 - (r_1 \overline{r_2} + \overline{r_1} r_2) \\ &= 2 \left[\sum_{j=0}^3 r_{1j}^2 + (\xi + \xi^4) \sum_{j=1}^3 r_{1j} r_{1,j-1} + (\xi^2 + \xi^3) \sum_{j-k \geq 2} r_{1j} r_{1k} \right] \\ &+ 2 \left[\sum_{j=0}^3 r_{2j}^2 + (\xi + \xi^4) \sum_{j=1}^3 r_{2j} r_{2,j-1} + (\xi^2 + \xi^3) \sum_{j-k \geq 2} r_{2j} r_{2k} \right] \\ &- \left[2 \sum_{j=0}^3 r_{1j} r_{2j} + (\xi + \xi^4) \sum_{j=1}^3 (r_{1j} r_{2,j-1} + r_{1,j-1} r_{2j}) \right. \\ &\quad \left. + (\xi^2 + \xi^3) \sum_{j-k \geq 2} (r_{1j} r_{2k} + r_{1k} r_{2j}) \right] \\ &= A_1 + A_2 \cdot \frac{\sqrt{5}}{2} \quad \text{with } A_1, A_2 \in \mathbb{Z}. \end{aligned}$$

It is not so easy to find, but easy to check that A_1 is equal to

$$\begin{aligned} &\frac{1}{4} \sum_{j=0}^3 [r_{1j}^2 + r_{2j}^2 + (r_{1j} - r_{2j})^2] \\ &+ \frac{1}{4} \sum_{j < k} [(r_{1j} - r_{1k} - r_{2j} + r_{2k})^2 + (r_{1j} - r_{1k})^2 + (r_{2j} - r_{2k})^2]. \end{aligned} \quad (56)$$

Now it is an easy exercise to find the 8-tuples $(r_{10}, \dots, r_{23}) \in \mathbb{Z}^8$ for which (56) takes the value 2. They are (here $e_j = (\delta_{ij})_{i=1, \dots, 8}$ for $j = 1, \dots, 8$ is the standard basis of $\mathbb{Z}^8$)

$$\begin{aligned} &\pm e_1, \dots, \pm e_8, \pm(e_1 + e_5), \pm(e_2 + e_6), \pm(e_3 + e_7), \pm(e_4 + e_8), \\ &\pm(1, 1, 1, 1, 0, 0, 0, 0), \pm(0, 0, 0, 0, 1, 1, 1, 1), \pm(1, 1, 1, 1, 1, 1, 1, 1). \end{aligned} \quad (57)$$

Observe $1 + \xi + \xi^2 + \xi^3 = -\xi^4$ and

$$\{r \in V_{\mathbb{Z}} \mid L_{\mathbb{Z}}(r, r) = 2\} = \{\pm b_1, \pm b_2, \pm(b_1 + b_2)\}.$$

The coefficients $(r_{10}, \dots, r_{23})$ in (57) give precisely the elements $r = r_1 b_1 + r_2 b_2$ on the right hand side of (46). This shows part (a) for $(m, l) = (2, 5)$.

(b) Any element g of $\text{Aut}(V_{\mathbb{Z}[\xi]}, L_{\mathbb{C}})$ will map the sets in (46) and (47) to themselves. The basis elements b_1 and b_2 are mapped to two elements in these sets with $L_{\mathbb{C}}(g(b_1), g(b_2)) = L_{\mathbb{Z}}(b_1, b_2) = -1$. Therefore g is up to a factor in $\{\pm \xi^k \mid k \in \mathbb{Z}\}$ an element of $\text{Aut}(V_{\mathbb{Z}}, L_{\mathbb{Z}})$. This shows (48).

In the case $m \geq 3$

$$\begin{aligned} \{r \in V_{\mathbb{Z}} \mid L_{\mathbb{Z}}(r, r) = 2\} &= \{\pm b_1\} \\ \text{and } \{r \in V_{\mathbb{Z}} \mid L_{\mathbb{Z}}(r, r) = m\} &= \{\pm b_2, \pm(b_1 + b_2)\}. \end{aligned}$$

This shows (49). The case $m = 2$ is the case of the root lattice of type A_2 . (50) is well known and easy to see.

(c) In the cases Z_{12} and Z_{18} as curve singularities, the examples 2.2 (ii) and (iii) showed $\text{Aut}(\text{Rad}(I), L) = \{\pm \text{id}\}$. Here $\text{Rad}(I) = \ker(M_h^{\text{curve case}} - \text{id})$. Under stabilization $\text{Rad}(I)$ becomes B_3 , and L changes just the sign, see (16). Thus $\text{Aut}(B_3, L) = \{\pm \text{id}\}$. Because of $M_h|_{B_3} = -\text{id}$ and $U := \{\text{id}\}$ in the cases Z_{12} and Z_{18} , this shows (c) in these cases.

Now consider the cases Q_{12}, Q_{16}, U_{12} and U_{16} . Here part (b) will be used, but that has to be prepared.

The normal forms of the quasihomogeneous surface singularities, given in section 7, show that they are sums of singularities in different variables of types A_l and D_{2m} with $(l, 2m)$ as follows,

	Q_{12}	Q_{16}	U_{12}	U_{16}	
$(l, 2m)$	$(2, 6)$	$(2, 8)$	$(3, 4)$	$(4, 4)$	(58)
	$A_2 \otimes D_6$	$A_2 \otimes D_8$	$A_3 \otimes D_4$	$A_4 \otimes D_4$	

Here the singularity A_l is in one variable and has the characteristic polynomial $p_{ch}^{A_l} = (t^{l+1} - 1)/(t - 1)$, the singularity D_{2m} is a curve singularity and has the characteristic polynomial $p_{ch}^{D_{2m}} = (t^{2m-1} - 1)\Phi_1$. The Thom-Sebastiani results which were cited in section 2 apply,

$$\begin{aligned} (Ml(f), L) &\cong (Ml(A_l), L_{A_l}) \otimes (Ml(D_{2m}), L_{D_{2m}}), \\ M_h &\cong M_h^{A_l} \otimes M_h^{D_{2m}}, \end{aligned} \quad (59)$$

and show

$$\begin{aligned} p_2 &= p_{ch}^{A_l}, \\ (B_3, L) &\cong (Ml(A_l), L_{A_l}) \otimes (Ml(D_{2m})_{1, \mathbb{Z}}, L_{D_{2m}}), \\ M_h|_{B_3} &\cong M_h^{A_l} \otimes \text{id}. \end{aligned} \quad (60)$$

The pair $(Ml(D_{2m})_{1,\mathbb{Z}}, L_{D_{2m}})$ was considered in example 2.2 (i). There is a $\mathbb{Z}$ -basis $\underline{b} = (b_1, b_2)$ of $Ml(D_{2m})_{1,\mathbb{Z}}$ with

$$L_{D_{2m}}(\underline{b}^t, \underline{b}) = \begin{pmatrix} -2 & 1 \\ 1 & -m \end{pmatrix}. \quad (61)$$

The pairings L and $L_{A_l} \otimes L_{D_{2m}}$ will be extended sesquilinearly from the $\mathbb{Z}$ -lattices to the $\mathbb{C}$ -vector spaces.

The $\mathbb{Z}$ -lattice $Ml(A_l)$ is a cyclic monodromy module. Choose a generator e of it. Therefore $Ml(A_l) \otimes Ml(D_{2m})_{1,\mathbb{Z}}$ is a sum of two cyclic monodromy modules, and generators are $e \otimes b_1$ and $e \otimes b_2$. For any automorphism g of $(Ml(A_l) \otimes Ml(D_{2m})_{1,\mathbb{Z}}, M_h^{A_l} \otimes \text{id})$ there are unique polynomials $g_1, g_2, g_3, g_4 \in \mathbb{Z}[t]$ of degree $\leq \deg p_2 - 1$ such that

$$\begin{pmatrix} g(v \otimes b_1) \\ g(v \otimes b_2) \end{pmatrix} = \begin{pmatrix} g_1(M_h^{A_l})(v) \otimes b_1 + g_3(M_h^{A_l})(v) \otimes b_2 \\ g_2(M_h^{A_l})(v) \otimes b_1 + g_4(M_h^{A_l})(v) \otimes b_2 \end{pmatrix} \quad (62)$$

for any $v \in Ml(A_l)$.

Now choose any eigenvalue ξ of $M_h^{A_l}$. Then $\mathbb{Z}[\xi]$ is a principal ideal domain. The space $\ker(M_h^{A_l} - \xi \text{id}) \cap Ml(A_l)_{\mathbb{Z}[\xi]}$ is a free $\mathbb{Z}[\xi]$ -module of rank 1. Choose a generating vector v . This choice gives an isomorphism from this space to $\mathbb{Z}[\xi]$. The spaces

$\ker(M_h - \xi \text{id}) \cap Ml(f)_{\mathbb{Z}[\xi]} \cong (\ker(M_h^{A_l} - \xi \text{id}) \cap Ml(A_l)_{\mathbb{Z}[\xi]}) \otimes Ml(D_{2m})_{1,\mathbb{Z}[\xi]}$ are free $\mathbb{Z}[\xi]$ -modules of rank 2. The space on the right hand side has the $\mathbb{Z}[\xi]$ -basis $(v \otimes b_1, v \otimes b_2) =: v \otimes \underline{b}$. Now (62) becomes

$$g(v \otimes \underline{b}) = v \otimes \underline{b} \cdot \begin{pmatrix} g_1(\xi) & g_2(\xi) \\ g_3(\xi) & g_4(\xi) \end{pmatrix}. \quad (63)$$

The pairing satisfies

$$(L_{A_l} \otimes L_{D_{2m}})(v \otimes \underline{b}) = L_{A_l}(v, v) \cdot \begin{pmatrix} -2 & 1 \\ 1 & -m \end{pmatrix}, \quad (64)$$

where $L_{A_l}(v, v) \in \mathbb{Z}[\xi] \cap \mathbb{R}_{>0}$. This space with this pairing is up to a scalar isomorphic to a pair $(V_{\mathbb{Z}[\xi]}, L_{\mathbb{C}})$ considered in the parts (a) and (b). Therefore by part (b), its group of automorphisms is isomorphic to $\{\pm \xi^k \mid k \in \mathbb{Z}\} \cdot \text{Aut}(V_{\mathbb{Z}}, L_{\mathbb{Z}})$.

Thus any element of $\text{Aut}(B_3, L)$ restricts on $\ker(M_h - \xi \text{id}) \cap Ml(f)_{\mathbb{Z}[\xi]}$ to such an automorphism. In the cases Q_{12}, Q_{16} and U_{16} the polynomial $p_2 = \Phi_3, \Phi_3$ respectively Φ_5 is irreducible, so all its zeros ξ are conjugate. Therefore then

$$\begin{aligned} \text{Aut}(B_3, L) &\cong \{\pm (M_h|_{B_3})^k \mid k \in \mathbb{Z}\} \cdot \text{Aut}(V_{\mathbb{Z}}, L_{\mathbb{Z}}) \\ &\cong \{\pm (M_h|_{B_3})^k \mid k \in \mathbb{Z}\} \times U, \end{aligned}$$

which proves (c) in these cases.

In the case U_{12} the characteristic polynomial $p_{ch}^{A_3} = p_2 = \Phi_4 \Phi_2$ is reducible. Consider an automorphism g of

$$(Ml(A_l) \otimes Ml(D_{2m})_{1,\mathbb{Z}}, L_{A_l} \otimes L_{D_{2m}}).$$

It is determined by the polynomials $g_1, g_2, g_3, g_4 \in \mathbb{Z}[t]$ in (62). For $\xi = i$ and for $\xi = -1$ it gives an automorphism of $\mathbb{Z}[\xi]v \otimes b_1 \oplus \mathbb{Z}[\xi]v \otimes b_2$ which is given by a matrix $\begin{pmatrix} g_1(\xi) & g_2(\xi) \\ g_3(\xi) & g_4(\xi) \end{pmatrix}$ which is by part (b) in

$$\{\pm \xi^k \mid k \in \mathbb{Z}\} \cdot \left\{ \pm \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \pm \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}, \pm \begin{pmatrix} -1 & 1 \\ -1 & 0 \end{pmatrix}, \right. \quad (65) \\ \left. \pm \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \pm \begin{pmatrix} -1 & 0 \\ -1 & 1 \end{pmatrix}, \pm \begin{pmatrix} 1 & -1 \\ 0 & -1 \end{pmatrix} \right\}.$$

By multiplying g with a suitable automorphism we can suppose that the matrix for $\xi = i$ is the identity matrix. Then

$$\begin{pmatrix} g_1 & g_2 \\ g_3 & g_4 \end{pmatrix} = \begin{pmatrix} 1 + (t^2 + 1)\tilde{g}_1 & (t^2 + 1)\tilde{g}_2 \\ (t^2 + 1)\tilde{g}_3 & 1 + (t^2 + 1)\tilde{g}_4 \end{pmatrix},$$

for some $\tilde{g}_1, \tilde{g}_2, \tilde{g}_3, \tilde{g}_4 \in \mathbb{Z}[t]$, so

$$\begin{pmatrix} g_1(-1) & g_2(-1) \\ g_3(-1) & g_4(-1) \end{pmatrix} = \begin{pmatrix} 1 + 2\tilde{g}_1(-1) & 2\tilde{g}_2(-1) \\ 2\tilde{g}_3(-1) & 1 + 2\tilde{g}_4(-1) \end{pmatrix}.$$

The only two possibilities are $\pm \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. In the case of a minus sign $g \circ ((M_h^{A_l})^2 \otimes \text{id}_{D_{2m}}) = -\text{id}$, in the case of a plus sign $g = \text{id}$. This finishes the proof of (c) in the case U_{12} .

(d) In section 7 a subgroup G^{mar} of $G_{\mathbb{Z}}$ will be calculated, and it will be shown that $G^{\text{mar}} = \{\pm M_h^k \mid k \in \mathbb{Z}\} \times U$. This shows $G_{\mathbb{Z}} \supset \{\pm M_h^k \mid k \in \mathbb{Z}\} \times U$ and that the map $G_{\mathbb{Z}} \rightarrow \text{Aut}(B_3, L)$ is surjectiv. $\square$

Remark 4.3. The number theoretic fact $\mathbb{Z}[e^{2\pi i/a}] \cap S^1 = \{\pm e^{2\pi i k/a} \mid k \in \mathbb{Z}\}$ can be interpreted as saying that in the case of the A_1 -lattice $V_{\mathbb{Z}} = \mathbb{Z}$ with $\mathbb{Z}$ -basis $b_1 = 1$ and standard bilinear form $L_{\mathbb{Z}}$ with $L_{\mathbb{Z}}(b_1, b_1) = 1$ and hermitian extension $L_{\mathbb{C}}$ to $\mathbb{C}$ the analogue of (46) holds. Now (46) for $m = 2$ can be seen as a generalization from the case A_1 to the case A_2 . Above it is proved only in the cases $l = 3, 4, 5$.

5. REVIEW ON μ -CONSTANT MONODROMY GROUPS G^{mar} , MARKED SINGULARITIES, THEIR MODULI SPACES M_{μ}^{mar} , AND TORELLI TYPE CONJECTURES

This paper is a sequel to [He7]. That paper studied holomorphic functions germs $f : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$ with an isolated singularity at 0

from a global perspective. Here we review most of the notions and results from [He7].

It defined the notions of *marked singularity* and *strongly marked singularity*. The marking uses the Milnor lattice $Ml(f) \cong \mathbb{Z}^\mu$ and the Seifert form L on it, which are explained in section 2.

Definition 5.1. Fix one reference singularity f_0 .

(a) Then a strong marking for any singularity f in the μ -homotopy class of f_0 (i.e. there is a 1-parameter family of singularities with constant Milnor number connecting f and f_0) is an isomorphism $\rho : (Ml(f), L) \rightarrow (Ml(f_0), L)$.

(b) The pair (f, ρ) is a *strongly marked singularity*. Two strongly marked singularities (f_1, ρ_1) and (f_2, ρ_2) are right equivalent (notation: $\sim_{\mathcal{R}}$) if a coordinate change $\varphi : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}^{n+1}, 0)$ with

$$f_1 = f_2 \circ \varphi \quad \text{and} \quad \rho_1 = \rho_2 \circ \varphi_{hom}$$

exists, where $\varphi_{hom} : (Ml(f_1), L) \rightarrow (Ml(f_2), L)$ is the induced isomorphism.

(c) The notion of a marked singularity is slightly weaker. If f and ρ are as above, then the pair $(f, \pm\rho)$ is a *marked singularity* (writing $\pm\rho$, the set $\{\rho, -\rho\}$ is meant, neither ρ nor $-\rho$ is preferred).

(d) Two marked singularities (f_1, ρ_1) and (f_2, ρ_2) are right equivalent (notation: $\sim_{\mathcal{R}}$) if a coordinate change φ with

$$f_1 = f_2 \circ \varphi \quad \text{and} \quad \rho_1 = \pm\rho_2 \circ \varphi_{hom}$$

exists.

Remarks 5.2. (i) The notion of a marked singularity behaves better than the notion of a strongly marked singularity, because it is not known whether all μ -homotopy families of singularities satisfy one of the following two properties:

Assumption (5.1): Any singularity in the μ -homotopy class of f_0 has multiplicity ≥ 3 . (66)

Assumption (5.2): Any singularity in the μ -homotopy class of f_0 has multiplicity 2. (67)

We expect that always one of two assumptions holds. For curve singularities and singularities right equivalent to semiquasihomogeneous singularities this is true, but in general it is not known. In a μ -homotopy family where neither of the two assumptions holds, strong marking behaves badly, see (ii).

(ii) If $\text{mult}(f) = 2$ then $(f, \rho) \sim_{\mathcal{R}} (f, -\rho)$, which is easy to see. If $\text{mult}(f) \geq 3$, then $(f, \rho) \not\sim_{\mathcal{R}} (f, -\rho)$, whose proof in [He7] is quite

intricate. These properties imply that the moduli space for strongly marked singularities discussed below is not Hausdorff in the case of a μ -homotopy class which satisfies neither one of the assumptions (66) or (67)

In [He6] for the μ -homotopy class of any singularity f_0 a moduli space $M_\mu(f_0)$ was constructed. As a set it is simply the set of right equivalence classes of singularities in the μ -homotopy class of f_0 . But in [He6] it is constructed as an analytic geometric quotient, and it is shown that it is locally isomorphic to the μ -constant stratum of a singularity modulo the action of a finite group. The μ -constant stratum of a singularity is the germ $(S_\mu, 0) \subset (M, 0)$ within the germ of the base space of a universal unfolding F of f , such that for a suitable representative

$$S_\mu = \{t \in M \mid F_t \text{ has only one singularity } x_0 \text{ and } F_t(x_0) = 0\}. \quad (68)$$

It comes equipped with a canonical complex structure, and M_μ inherits a canonical structure, see the chapters 12 and 13 in [He6].

In [He7] analogous results for marked singularities were proved. A better property is that M_μ^{mar} is locally isomorphic to a μ -constant stratum without dividing out a finite group action. Therefore one can consider it as a *global μ -constant stratum* or as a *Teichmüller space for singularities*. The following theorem collects results from [He7, theorem 4.3].

Theorem 5.3. *Fix one reference singularity f_0 . Define the sets*

$$M_\mu^{smar}(f_0) := \{\text{strongly marked } (f, \rho) \mid f \text{ in the } \mu\text{-homotopy class of } f_0\} / \sim_{\mathcal{R}}, \quad (69)$$

$$M_\mu^{mar}(f_0) := \{\text{marked } (f, \pm\rho) \mid f \text{ in the } \mu\text{-homotopy class of } f_0\} / \sim_{\mathcal{R}}. \quad (70)$$

(a) $M_\mu^{mar}(f_0)$ carries a natural canonical complex structure. It can be constructed with the underlying reduced complex structure as an analytic geometric quotient (see [He7, theorem 4.3] for details).

(b) The germ $(M_\mu^{mar}(f_0), [(f, \pm\rho)])$ with its canonical complex structure is isomorphic to the μ -constant stratum of f with its canonical complex structure (see [He6, chapter 12] for the definition of that).

(c) For any $\psi \in G_{\mathbb{Z}}(f_0) =: G_{\mathbb{Z}}$, the map

$$\psi_{mar} : M_\mu^{mar} \rightarrow M_\mu^{mar}, \quad [(f, \pm\rho)] \rightarrow [(f, \pm\psi \circ \rho)]$$

is an automorphism of M_μ^{mar} . The action

$$G_{\mathbb{Z}} \times M_\mu^{mar} \rightarrow M_\mu^{mar}, \quad (\psi, [(f, \pm\rho)]) \mapsto \psi_{mar}([(f, \pm\rho)])$$

is a group action from the left.

(d) The action of $G_{\mathbb{Z}}$ on M_{μ}^{mar} is properly discontinuous. The quotient $M_{\mu}^{mar}/G_{\mathbb{Z}}$ is the moduli space M_{μ} for right equivalence classes in the μ -homotopy class of f_0 , with its canonical complex structure. Especially, $[(f_1, \pm\rho_1)]$ and $[(f_2, \pm\rho_2)]$ are in one $G_{\mathbb{Z}}$ -orbit if and only if f_1 and f_2 are right equivalent.

(e) If assumption (66) or (67) holds then (a) to (d) are also true for M_{μ}^{smar} and ψ_{smar} with $\psi_{smar}([(f, \rho)]) := [(f, \psi \circ \rho)]$. If neither (66) nor (67) holds then the natural topology on M_{μ}^{smar} is not Hausdorff.

We stick to the situation in theorem 5.3 and define two subgroups of $G_{\mathbb{Z}}(f_0)$. The definitions in [He7, definition 3.1] are different, they use μ -constant families. The following definitions are a part of theorem 4.4 in [He7].

Definition 5.4. Let $(M_{\mu}^{mar})^0$ be the topological component of M_{μ}^{mar} (with its reduced complex structure) which contains $[(f_0, \pm \text{id})]$. Then

$$G^{mar}(f_0) := \{\psi \in G_{\mathbb{Z}} \mid \psi \text{ maps } (M_{\mu}^{mar})^0 \text{ to itself}\} \subset G_{\mathbb{Z}}(f_0). \quad (71)$$

If assumption (66) or (67) holds, $(M_{\mu}^{smar})^0$ and $G^{smar}(f_0) \subset G_{\mathbb{Z}}(f_0)$ are defined analogously.

The following theorem is also proved in [He7].

Theorem 5.5. (a) In the situation above the map

$$\begin{aligned} G_{\mathbb{Z}}/G^{mar}(f_0) &\rightarrow \{\text{topological components of } M_{\mu}^{mar}\} \\ \psi \cdot G^{mar}(f_0) &\mapsto \text{the component } \psi_{mar}((M_{\mu}^{mar})^0) \end{aligned}$$

is a bijection.

(b) If assumption (66) or (67) holds then (a) is also true for M_{μ}^{smar} and $G^{smar}(f_0)$.

(c) $-\text{id} \in G_{\mathbb{Z}}$ acts trivially on $M_{\mu}^{mar}(f_0)$. Suppose additionally that assumption (66) holds for f_0 . Then $\{\pm \text{id}\}$ acts freely on $M_{\mu}^{smar}(f_0)$, and the quotient map

$$M_{\mu}^{smar}(f_0) \xrightarrow{/\{\pm \text{id}\}} M_{\mu}^{mar}(f_0), \quad [(f, \rho)] \mapsto [(f, \pm \rho)]$$

is a double covering.

The first main conjecture in [He7] is part (a) of the following conjecture (the second main conjecture in [He7] is conjecture 5.11 (a) below).

Conjecture 5.6. (a) Fix a singularity f_0 . Then M_{μ}^{mar} is connected. Equivalently (in view of theorem 5.5 (a)): $G^{mar}(f_0) = G_{\mathbb{Z}}$.

(b) If the μ -homotopy class of f_0 satisfies assumption (66), then $-\text{id} \notin G^{smar}(f_0)$.

If (a) holds then (b) is equivalent to $M_\mu^{smar}(f_0)$ having two components. If (a) and (b) hold, there should be a natural invariant which distinguishes the index 2 subgroup $G^{smar} \subset G^{mar} = G_{\mathbb{Z}}$. Anyway, part (a) is the more important conjecture. Using the other definition of G^{mar} in [He7], it says that up to $\pm \text{id}$, any element of $G_{\mathbb{Z}}$ can be realized as transversal monodromy of a μ -constant family with parameter space S^1 .

The whole conjecture 5.6 had been proved in [He7] for the simple singularities and those 22 of the 28 exceptional unimodal and bimodal singularities, where all eigenvalues of the monodromy have only multiplicity one [He7, theorems 8.3 and 8.4]. In this paper it will be proved for the remaining unimodal and exceptional bimodal singularities.

In order to understand the stabilizers $\text{Stab}_{G_{\mathbb{Z}}}([(f, \rho)])$ and $\text{Stab}_{G_{\mathbb{Z}}}([(f, \pm \rho)])$ of points $[(f, \rho)] \in M_\mu^{smar}(f_0)$ and $[(f, \pm \rho)] \in M_\mu^{mar}(f_0)$, we have to look at the *symmetries* of a single singularity. These had been discussed in [He6, chapter 13.2]. The discussion had been taken up again in [He7].

Definition 5.7. Let $f_0 = f_0(x_0, \dots, x_n)$ be a reference singularity and let f be any singularity in the μ -homotopy class of f_0 . If ρ is a marking, then $G_{\mathbb{Z}}(f) = \rho^{-1} \circ G_{\mathbb{Z}} \circ \rho$.

We define

$$\mathcal{R} := \{\varphi : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}^{n+1}, 0) \text{ biholomorphic}\}, \quad (72)$$

$$\mathcal{R}^f := \{\varphi \in \mathcal{R} \mid f \circ \varphi = f\}, \quad (73)$$

$$R_f := j_1 \mathcal{R}^f / (j_1 \mathcal{R}^f)^0, \quad (74)$$

$$G_{\mathcal{R}}^{smar}(f) := \{\varphi_{hom} \mid \varphi \in \mathcal{R}^f\} \subset G_{\mathbb{Z}}(f), \quad (75)$$

$$G_{\mathcal{R}}^{mar}(f) := \{\pm \psi \mid \psi \in G_{\mathcal{R}}^{smar}(f)\}. \quad (76)$$

Again, the definition of $G_{\mathcal{R}}^{smar}$ is different from the definition in [He7, definition 3.1]. The characterization in (75) is [He7, theorem 3.3. (e)]. R_f is the finite group of components of the group $j_1 \mathcal{R}^f$ of 1-jets of coordinate changes which leave f invariant. The following theorem collects results from several theorems in [He7].

Theorem 5.8. *Consider the data in definition 5.7.*

- (a) *If $\text{mult}(f) \geq 3$ then $j_1 \mathcal{R}^f = R_f$.*
- (b) *The homomorphism $()_{hom} : \mathcal{R}^f \rightarrow G_{\mathbb{Z}}(f)$ factors through R_f . Its image is $(R_f)_{hom} = G_{\mathcal{R}}^{smar}(f) \subset G_{\mathbb{Z}}(f)$.*
- (c) *The homomorphism $()_{hom} : R_f \rightarrow G_{\mathcal{R}}^{smar}(f)$ is an isomorphism.*
- (d)

$$-\text{id} \notin G_{\mathcal{R}}^{smar}(f) \iff \text{mult } f \geq 3. \quad (77)$$

Equivalently: $G_{\mathcal{R}}^{\text{mar}}(f) = G_{\mathcal{R}}^{\text{smar}}(f)$ if $\text{mult } f = 2$, and $G_{\mathcal{R}}^{\text{mar}}(f) = G_{\mathcal{R}}^{\text{smar}}(f) \times \{\pm \text{id}\}$ if $\text{mult } f \geq 3$.

(e) $G_{\mathcal{R}}^{\text{mar}}(f) = G_{\mathcal{R}}^{\text{mar}}(f + x_{n+1}^2)$.

(f) $M_h \in G^{\text{smar}}(f)$. If f is quasihomogeneous then $M_h \in G_{\mathcal{R}}^{\text{smar}}(f)$.

(g) For any $[(f, \rho)] \in M_{\mu}^{\text{smar}}$

$$\text{Stab}_{G_{\mathbb{Z}}}([(f, \rho)]) = \rho \circ G_{\mathcal{R}}^{\text{smar}}(f) \circ \rho^{-1}, \quad (78)$$

$$\text{Stab}_{G_{\mathbb{Z}}}([(f, \pm \rho)]) = \rho \circ G_{\mathcal{R}}^{\text{mar}}(f) \circ \rho^{-1}. \quad (79)$$

((78) does not require assumption (66) or (67)). As $G_{\mathbb{Z}}$ acts properly discontinuously on $M_{\mu}^{\text{mar}}(f_0)$, $G_{\mathcal{R}}^{\text{smar}}(f)$ and $G_{\mathcal{R}}^{\text{mar}}(f)$ are finite. (But this follows already from the finiteness of R_f and (b).)

In the case of a quasihomogeneous singularity the group R_f has a canonical lift to $\mathcal{R}^f$. It will be useful for the calculation of R_f .

Theorem 5.9. [He6, theorem 13.11] *Let $f \in \mathbb{C}[x_0, \dots, x_n]$ be a quasihomogeneous polynomial with an isolated singularity at 0 and weights $w_0, \dots, w_n \in \mathbb{Q} \cap (0, \frac{1}{2}]$ and weighted degree 1. Suppose that $w_0 \leq \dots \leq w_{n-1} < \frac{1}{2}$ (then $f \in \mathfrak{m}^3$ if and only if $w_n < \frac{1}{2}$). Let G_w be the algebraic group of quasihomogeneous coordinate changes, that means, those which respect $\mathbb{C}[x_0, \dots, x_n]$ and the grading by the weights $w_0, \dots, w_n$ on it. Then*

$$R_f \cong \text{Stab}_{G_w}(f). \quad (80)$$

Finally we need and we want to study period maps and Torelli type problems for singularities.

This story should start with the definition of the Gauß-Manin connection and the Brieskorn lattice for an isolated hypersurface singularity. This had been developed in many papers of the second author, and also much earlier by Brieskorn, K. Saito, G.-M. Greuel, F. Pham, A. Varchenko, M. Saito and others.

As we will build here on calculations done in [He1] and therefore never have to touch Brieskorn lattices explicitly, we take here a formal point of view and refer to [He1], [He2], [He4], [He6] and [He7] for the definitions of the following objects.

Any singularity f comes equipped with a *Brieskorn lattice* $H_0''(f)$. It is much richer than, but still comparable to a Hodge structure of a closed Kähler manifold.

After fixing a reference singularity f_0 , a marked singularity $(f, \pm \rho)$ comes equipped with a *marked Brieskorn lattice* $BL(f, \pm \rho)$. A classifying space $D_{BL}(f_0)$ for marked Brieskorn lattices was constructed in [He4]. It is especially a complex manifold, and $G_{\mathbb{Z}}$ acts properly discontinuously on it.

Theorem 5.10. *Fix one reference singularity f_0 .*

(a) *There is a natural holomorphic period map*

$$BL : M_\mu^{mar}(f_0) \rightarrow D_{BL}(f_0). \quad (81)$$

It is $G_{\mathbb{Z}}$ -equivariant.

(b) [He6, theorem 12.8] *It is an immersion, here the reduced complex structure on $M_\mu^{mar}(f_0)$ is considered. (The second author has also a proof that it is an immersion where the canonical complex structure on $M_\mu^{mar}(f_0)$ is considered, but the proof is not written).*

The second main conjecture in [He7] is part (a) of the following conjecture. Part (a) and part (b) are global Torelli type conjectures.

Conjecture 5.11. *Fix one reference singularity f_0 .*

(a) *The period map $BL : M_\mu^{mar} \rightarrow D_{BL}$ is injective.*

(b) *The period map $LBL : M_\mu = M_\mu^{mar}/G_{\mathbb{Z}} \rightarrow D_{BL}/G_{\mathbb{Z}}$ is injective.*

(c) *For any singularity f in the μ -homotopy class of f_0 and any marking ρ ,*

$$\text{Stab}_{G_{\mathbb{Z}}}([(f, \pm\rho)]) = \text{Stab}_{G_{\mathbb{Z}}}(BL([(f, \pm\rho)])) \quad (82)$$

(only $\subset$ and the finiteness of both groups are clear).

The second author has a long-going project on Torelli type conjectures. Already in [He1], part (b) was conjectured and proved for all simple and unimodal singularities and almost all bimodal singularities (all except 3 subseries of the 8 bimodal series). This was possible without the general construction of M_μ and D_{BL} , which came later in [He6] and [He4]. In the concrete cases considered in [He1], it is easy to identify a posteriori the spaces M_μ and D_{BL} . We will make use of that in the sections 6 and 7. Part (c) was conjectured in [He6].

The following lemma from [He7] clarifies the logic between the parts (a), (b) and (c) of conjecture 5.11.

Lemma 5.12. *In conjecture 5.11, (a) $\iff$ (b) and (c).*

Part (a) of conjecture 5.11 was proved in [He7] for the simple and those 22 of the 28 exceptional unimodal and bimodal singularities, where all eigenvalues of the monodromy have multiplicity one. Here it will be proved for the remaining unimodal and the remaining exceptional bimodal singularities.

As part (b) of conjecture 5.11 was already proved in all these cases in [He1], the main work in [He7, section 8] and here is the control of the group $G_{\mathbb{Z}}$. This is carried out here in the sections 3 and 4, and it is surprisingly difficult.

6. G^{mar} , M_μ^{mar} AND A STRONG TORELLI RESULT FOR THE SIMPLE ELLIPTIC AND THE HYPERBOLIC SINGULARITIES

The 1-parameter families of the hyperbolic singularities of type T_{pqr} ($p, q, r \in \mathbb{N}_{\geq 2}$, $p \geq q \geq r$, $\kappa := \frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$) have as surface singularities the normal forms [AGV1]

$$x^p + y^q + z^r + t \cdot xyz, \quad t \in X := \mathbb{C}^*. \quad (83)$$

The 1-parameter families of the simple elliptic singularities $T_{333} = \tilde{E}_6$, $T_{442} = \tilde{E}_7$, $T_{632} = \tilde{E}_8$ have as surface singularities different normal forms [SaK]. The normal form $x^p + y^q + z^r + t \cdot xyz$ does in the case of T_{442} not contain representatives of all right equivalence classes, the class with j -invariant $j = 1$ is missing [SaK, 1.11, Bem. (ii)]. Therefore we work in the following also with the Legendre normal forms

$$\begin{aligned} T_{333} : \quad & y(y-x)(y-\lambda x) - xz^2, \quad t \in X := \mathbb{C} - \{0, 1\}, \\ T_{442} : \quad & yx(y-x)(y-\lambda x) + z^2, \quad t \in X := \mathbb{C} - \{0, 1\}, \\ T_{632} : \quad & y(y-x^2)(y-\lambda x^2) + z^2, \quad t \in X := \mathbb{C} - \{0, 1\}. \end{aligned} \quad (84)$$

They contain representatives of all right equivalence classes. Let X^{univ} denote the universal covering of X , so $X^{univ} = \mathbb{C}$ if $\kappa < 1$ and $X^{univ} = \mathbb{H}$ if $\kappa = 1$.

Theorem 6.1. (a) *For the simple elliptic singularities and the hyperbolic singularities in any number of variables, the space M_μ^{mar} of right equivalence classes of marked singularities is $M_\mu^{mar} \cong X^{univ}$, so it is connected, and thus $G^{mar} = G_{\mathbb{Z}}$. The period map $BL : M_\mu^{mar} \rightarrow D_{BL}$ is an isomorphism, so the strong global Torelli conjecture 5.11 (a) is true.*

(b) *Now consider the singularities of type T_{pqr} as curve singularities if $r = 2$ and as surface singularities if $r \geq 3$. Then*

$$G_{\mathbb{Z}} = G^{mar} = G^{smar} \times \{\pm \text{id}\}, \quad \text{equivalently: } -\text{id} \notin G^{smar}. \quad (85)$$

The subgroup of G^{smar} , which acts trivially on M_μ^{mar} , is the kernel of the surjective map

$$G^{smar} \rightarrow \text{Aut}(Ml(f_0)_{1,\mathbb{Z}}, L)/\{\pm \text{id}\}. \quad (86)$$

It is equal to $\rho \circ (\mathcal{R}^f)_{\text{hom}} \circ \rho^{-1}$ for a generic $[(f, \rho)] \in M_\mu^{mar}$. Its size is 54, 16 and 6 for T_{333} , T_{442} and T_{632} .

Proof: (a) The proof uses two Torelli type results from [He1].

We choose a marked reference singularity $[(f_0, \pm \text{id})]$ in X^{univ} , then all elements of X^{univ} become marked singularities, because X^{univ} is simply connected. Then the period map $X^{univ} \rightarrow D_{BL}$ is well defined.

The first Torelli type result from [He1] is that this map is an isomorphism.

Therefore the marked Brieskorn lattices of the marked singularities in X^{univ} are all different. Therefore the marked singularities in X^{univ} are all not right equivalent. This gives an embedding $X^{univ} \hookrightarrow M_\mu^{mar}(f_0)^0$.

On the other hand, we have the period map $BL : M_\mu^{mar}(f_0)^0 \rightarrow D_{BL}$, which is an immersion. As it restricts to the isomorphism $X^{univ} \rightarrow D_{BL}$, finally $X^{univ} = M_\mu^{mar}(f_0)^0$.

For part (a) it rests to show that $G^{mar} = G_{\mathbb{Z}}$. Then M_μ^{mar} is connected and $M_\mu^{mar} = X^{univ}$, and $BL : M_\mu^{mar} \rightarrow D_{BL}$ is an isomorphism.

We have to look closer at D_{BL} and the action of $G_{\mathbb{Z}}$ on it. In the case $\kappa < 1$,

$$D_{BL} \cong \{V \subset Ml(f_0)_1 \mid \dim V = 1, V \neq \ker(M_h - \text{id})\}. \quad (87)$$

In the case $\kappa = 1$,

$$D_{BL} \cong \text{one component of } \{V \subset Ml(f_0)_1 \mid \dim V = 1, V \neq \bar{V}\}. \quad (88)$$

In both cases the group $\text{Aut}(Ml(f_0)_{1,\mathbb{Z}}, L)/\{\pm \text{id}\}$ acts faithfully on D_{BL} .

In both cases the second Torelli type result from [He1] which we need is that the period map

$$X^{univ} / \sim_R \rightarrow D_{BL} / \text{Aut}(Ml(f_0)_{1,\mathbb{Z}}, L) \quad (89)$$

is an isomorphism. Here $\sim_R$ denotes right equivalence for unmarked singularities.

Because of $X^{univ} = M_\mu^{mar}(f_0)^0$,

$$X^{univ} / \sim_R = M_\mu^{mar}(f_0)^0 / G^{mar}. \quad (90)$$

The isomorphism (89) and the isomorphism $M_\mu^{mar}(f_0)^0 = X^{univ} \rightarrow D_{BL}$ show that the map

$$G^{mar} \rightarrow \text{Aut}(Ml(f_0)_{1,\mathbb{Z}}, L) \quad (91)$$

is surjective. This completes the proof of part (b) of theorem 3.1.

It also shows that for proving $G^{mar} = G_{\mathbb{Z}}$, it is sufficient to show that the kernels of the maps to $\text{Aut}(Ml(f_0)_{1,\mathbb{Z}}, L)/\{\pm \text{id}\}$ coincide. The kernel of the map $G_{\mathbb{Z}} \rightarrow \text{Aut}(Ml(f_0)_{1,\mathbb{Z}}, L)/\{\pm \text{id}\}$ was determined in theorem 3.1. In fact, this is the only part of theorem 3.1 which we need here. It consists of those elements of $(U_1 \rtimes U_2) \times \{\pm \text{id}\}$ in (22) for which $\delta = 0$, so it is isomorphic to the group

$$\begin{aligned} & \left(\{(\alpha, \beta, \gamma) \in \mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r \mid \frac{\alpha}{p} + \frac{\beta}{q} + \frac{\gamma}{r} \equiv 0 \pmod{1}\} \rtimes U_2 \right) \times \{\pm \text{id}\} \\ & =: (U_1^0 \rtimes U_2) \times \{\pm \text{id}\}. \end{aligned} \quad (92)$$

The kernel of the map $G^{mar} \rightarrow \text{Aut}(Ml(f_0)_{1,\mathbb{Z}}, L)/\{\pm \text{id}\}$ is the subgroup of G^{mar} which acts trivially on $M_\mu^{mar}(f_0)^0$. It is the isotropy group in G^{mar} of a generic point $[(f, \pm \rho)] \in M_\mu^{mar}(f_0)^0$. So by theorem 5.8 (g) it is the group

$$\rho \circ G_{\mathcal{R}}^{mar}(f) \circ \rho^{-1} = \rho \circ \{\pm \varphi_{hom} \mid \varphi \in \mathcal{R}^f\} \circ \rho^{-1}. \quad (93)$$

As f is generic, we can and will use now the normal form $f = x^p + y^q + z^r + t \cdot xyz$, also in the case $\kappa = 1$. The following coordinate changes generate a finite subgroup $S \subset \mathcal{R}^f$

$$\begin{aligned} \varphi^{\alpha, \beta, \gamma} : (x, y, z) &\mapsto (e^{2\pi i \alpha/p} x, e^{2\pi i \beta/q} y, e^{2\pi i \gamma/r} z) \\ &\text{with } \frac{\alpha}{p} + \frac{\beta}{q} + \frac{\gamma}{r} \equiv 0 \pmod{1}, \\ \varphi^{1,2} : (x, y, z) &\mapsto (y, x, z) \quad \text{if } p = q, \\ \varphi^{2,3} : (x, y, z) &\mapsto (x, z, y) \quad \text{if } q = r, \\ \varphi^{minus} : (x, y, z) &\mapsto (x, y, -z - txy) \quad \text{if } r = 2. \end{aligned} \quad (94)$$

φ^{minus} has order 2 and commutes with the other coordinate changes ($q = r = 2$ is impossible because of $\kappa \leq 1$). The group S is isomorphic (as an abstract group) to $U_1^0 \rtimes U_2$ if $r \geq 3$ and to $(U_1^0 \rtimes U_2) \times \{\pm \text{id}\}$ if $r = 2$. The map to 1-jets of coordinate changes is injective,

$$S \xrightarrow{\cong} j_1 S \subset j_1 \mathcal{R}^f \subset j_1 \mathcal{R}. \quad (95)$$

Now we have to treat the cases $r \geq 3$ and $r = 2$ separately.

The case $r \geq 3$: Then $j_1 \mathcal{R}^f$ is finite and isomorphic to R_f , the map

$$(\cdot)_{hom} : R_f \rightarrow G_{\mathbb{Z}}(f) = \rho^{-1} \circ G_{\mathbb{Z}} \circ \rho$$

is injective, and the image $G_{\mathcal{R}}^{mar}(f)$ does not contain $-\text{id}$ (theorem 5.8). Therefore then $S \cong (S)_{hom} \subset G_{\mathbb{Z}}(f)$ and $-\text{id} \notin (S)_{hom}$. Thus the group $(S)_{hom} \times \{\pm \text{id}\}$ is isomorphic to $(U_1^0 \rtimes U_2) \times \{\pm \text{id}\}$. Now it is clear that the group in (93) is at least as big as the group in (92). But it cannot be bigger. So they are of equal size. This implies $G^{mar} = G_{\mathbb{Z}}$.

The case $r = 2$: We claim that the map $S \rightarrow (S)_{hom}$ is injective. If this is true then $(S)_{hom} \cong (U_1^0 \rtimes U_2) \times \{\pm \text{id}\}$, and this is of equal size as the group in (92). Then again the group in (93) is at least as big as the group in (92), but it cannot be bigger. So they are of equal size. This implies $G^{mar} = G_{\mathbb{Z}}$.

It rests to prove the claim. For this we consider the curve singularity

$$g := x^p + y^q - \frac{1}{4}tx^2y^2. \quad (96)$$

Then

$$\begin{aligned}
 g + z^2 &= f \circ \psi \quad \text{with } \psi(x, y, z) = (x, y, z - \frac{1}{2}txy), \\
 \mathcal{R}^{g+z^2} &= \psi^{-1} \circ \mathcal{R}^f \circ \psi, \\
 \psi^{-1} \circ \varphi^{\alpha, \beta, \gamma} \circ \psi &= \varphi^{\alpha, \beta, \gamma}, \\
 \psi^{-1} \circ \varphi^{1, 2} \circ \psi &= \varphi^{1, 2}, \quad \text{if } p = q, \\
 \psi^{-1} \circ \varphi^{minus} \circ \psi &= ((x, y, z) \mapsto (x, y, -z)).
 \end{aligned} \tag{97}$$

($q = r = 2$ is impossible because of $\kappa \leq 1$). The subgroup

$$S^{curve} := \{\varphi^{\alpha, \beta, \gamma} \circ (\varphi^{minus})^{-\gamma} \mid (\alpha, \beta, \gamma) \in U_1^0\} \rtimes U_2 \tag{98}$$

has index 2 in S , its conjugate $\psi^{-1} \circ S^{curve} \circ \psi$ restricts to $\mathcal{R}^g$, and it maps injectively to $j_1 \mathcal{R}^g \cong R_g$. By theorem 5.8 (c) the map $S^{curve} \rightarrow (S^{curve})_{hom}$ is injective, and $-\text{id}$ is not in the image. But $(\varphi^{minus})_{hom} = -\text{id}$. This proves the claim.

(b) By theorem 5.5 (c), the projection $M_\mu^{smar} \rightarrow M_\mu^{mar}$ is a twofold covering, and $-\text{id}$ exchanges the two sheets of the covering. Because of $M_\mu^{mar} = \mathbb{C}$ if $\kappa < 1$ and $M_\mu^{mar} = \mathbb{H}$ if $\kappa = 1$, M_μ^{smar} has two components. Therefore $-\text{id} \notin G^{smar}$ and $G_\mathbb{Z} = G^{mar} = G^{smar} \times \{\pm \text{id}\}$.

The statements right before and after (86) were already proved and used in the proof of part (a).

The group $\rho \circ (\mathcal{R}^f)_{hom} \circ \rho^{-1}$ for a generic $[(f, \rho)] \in M_\mu^{mar}$ has size 54, 16 and 6 for T_{333}, T_{442} and T_{632} , because it is isomorphic to an index 2 subgroup of the group in (92), and that group has 108, 32 and 12 elements in the cases T_{333}, T_{442} and T_{632} $\square$

7. G^{mar}, M_μ^{mar} AND A STRONG TORELLI RESULT FOR 6 OF THE 28 EXCEPTIONAL UNIMODAL AND BIMODAL SINGULARITIES

Normal forms for the 1-parameter families of the exceptional unimodal and bimodal singularities of types $Z_{12}, Q_{12}, U_{12}, Z_{18}, Q_{16}$ and U_{16} in the minimal number of variables are as follows [AGV1]. Here Z_{12} and Z_{18} are curve singularities, Q_{12}, U_{12}, Q_{16} and U_{16} are surface singularities. The singularity for the parameter $t = 0$ is quasihomogeneous, the others are semiquasihomogeneous. The space of the parameter $t = t_1$ or $t = (t_1, t_2)$ is $X = \mathbb{C}^{\text{mod}(f_0)}$. The weights $w = (w_x, w_y)$ respectively $w = (w_x, w_y, w_z)$ are normalized such that $\deg_w f_0 = 1$.

	normal form	$\text{mod}(f_0)$	weights
$Z_{12} :$	$f_t = x^3y + xy^4 + tx^2y^3$	1	$(\frac{3}{11}, \frac{2}{11})$
$Q_{12} :$	$f_t = x^3 + y^5 + yz^2 + txy^4$	1	$(\frac{1}{3}, \frac{1}{5}, \frac{2}{5})$
$U_{12} :$	$f_t = x^3 + y^3 + z^4 + txyz^2$	1	$(\frac{1}{3}, \frac{1}{3}, \frac{1}{4})$
$Z_{18} :$	$f_t = x^3y + xy^6 + (t_1 + t_2y)y^9$	2	$(\frac{3}{17}, \frac{2}{17})$
$Q_{16} :$	$f_t = x^3 + y^7 + yz^2 + (t_1 + t_2y)xy^5$	2	$(\frac{1}{3}, \frac{1}{7}, \frac{3}{7})$
$U_{16} :$	$f_t = x^3 + xz^2 + y^5 + (t_1 + t_2y)x^2y^2$	2	$(\frac{1}{3}, \frac{1}{5}, \frac{1}{3})$

(99)

The normal form of the quasihomogeneous singularity of type Q_{12} , Q_{16} , U_{12} and U_{16} is a sum of an A_l -singularity in one variable and a D_{2m} -singularity in two variables with $(l, 2m)$ as in table (100)=(58).

	Q_{12}	Q_{16}	U_{12}	U_{16}
$(l, 2m)$	$(2, 6)$	$(2, 8)$	$(3, 4)$	$(4, 4)$
	$A_2 \otimes D_6$	$A_2 \otimes D_8$	$A_3 \otimes D_4$	$A_4 \otimes D_4$

(100)

The rest of this section is devoted to the proof of the following theorem.

Theorem 7.1. (a) For the 6 families of exceptional unimodal and bimodal singularities of types $Z_{12}, Q_{12}, U_{12}, Z_{18}, Q_{16}$ and U_{16} in any number of variables, the space M_μ^{mar} of right equivalence classes of marked singularities is $M_\mu^{\text{mar}} \cong X = \mathbb{C}^{\text{mod}(f_0)}$, so it is connected, and thus $G^{\text{mar}} = G_{\mathbb{Z}}$. The period map $BL : M_\mu^{\text{mar}} \rightarrow D_{BL}$ is an isomorphism, so the strong global Torelli conjecture 5.11 (a) is true.

(b) Now consider the singularities of type Z_{12} and Z_{18} as curve singularities and the singularities of types Q_{12}, U_{12}, Q_{16} and U_{16} as surface singularities. Then their multiplicities are ≥ 3 . Then

$$G_{\mathbb{Z}} = G^{\text{mar}} = G^{\text{smar}} \times \{\pm \text{id}\}, \quad \text{equivalently: } -\text{id} \notin G^{\text{smar}}. \quad (101)$$

Proof: (a) The proof is similar to the proof of theorem 6.1, but simpler. We need only the first of the two Torelli type results from [He1], which were used in the proof of theorem 6.1.

We choose as marked reference singularity the quasihomogeneous singularity with trivial marking $[(f_0, \pm \text{id})]$ in X . Then all elements of X become marked singularities, because X is simply connected. Then the period map $X \rightarrow D_{BL}$ is well defined. A Torelli type result from [He1] says that this map is an isomorphism. It is in fact easy, because the singularities here are semiquasihomogeneous and only f_0 is quasihomogeneous. That makes the calculations easy.

Therefore the marked Brieskorn lattices of the marked singularities X are all different. Therefore the marked singularities in X are all not right equivalent. This gives an embedding $X \hookrightarrow M_\mu^{mar}(f_0)^0$.

On the other hand, we have the period map $BL : M_\mu^{mar}(f_0)^0 \rightarrow D_{BL}$, which is an immersion. As it restricts to the isomorphism $X \rightarrow D_{BL}$, finally $X = M_\mu^{mar}(f_0)^0$.

For part (a) it rests to show that $G^{mar} = G_{\mathbb{Z}}$. Then M_μ^{mar} is connected and $M_\mu^{mar} = X$, and $BL : M_\mu^{mar} \rightarrow D_{BL}$ is an isomorphism.

The weights of the deformation parameter(s) t_1 (and t_2) equip the parameter space $X = M_\mu^{mar}$ with a good $\mathbb{C}^*$ -action. It commutes with the action of $G_{\mathbb{Z}}$. This gives the first equality in

$$G^{mar} = \text{Stab}_{G_{\mathbb{Z}}}([(f_0, \pm \text{id})]) = G_{\mathcal{R}}^{mar}(f_0). \quad (102)$$

The second equality is part of theorem 5.8 (g).

On the other hand, by theorem 5.8 (d), $G_{\mathcal{R}}^{smar}(f_0) \times \{\pm \text{id}\} = G_{\mathcal{R}}^{mar}(f_0)$. But because f_0 is a quasihomogeneous singularity of degree ≥ 3 , the group $G_{\mathcal{R}}^{smar}(f_0)$ can be calculated easily via $\text{Stab}_{G_w}(f_0)$, see theorem 5.8 (c) and theorem 5.9: $\text{Stab}_{G_w}(f_0) \xrightarrow{\cong} G_{\mathcal{R}}^{smar}(f_0)$. Therefore it is sufficient to show that $\text{Stab}_{G_w}(f_0)$ has half as many elements as the group $G_{\mathbb{Z}}$. We postpone its proof. If it holds, then

$$G_{\mathbb{Z}} = G^{mar} = G_{\mathcal{R}}^{mar}(f_0) = G_{\mathcal{R}}^{smar}(f_0) \times \{\pm \text{id}\} \quad (103)$$

follows, and part (a) of the theorem is proved. For part (b), the same argument as in the proof of theorem 6.1 (b) works: M_μ^{smar} is a twofold covering of M_μ^{mar} , and the two sheets are exchanged by the action of $-\text{id}$. As $X = \mathbb{C}^{\text{mod}(f_0)}$, M_μ^{smar} has two components, and $-\text{id} \notin M_\mu^{smar}$.

In theorem 4.1 it was shown that $G_{\mathbb{Z}}$ is $G_{\mathbb{Z}} = \{\pm M_h^k \mid k \in \mathbb{Z}\} \times U$ with U as in table (39)=(104).

$$U \cong \begin{array}{c|c|c|c|c|c|c} & Z_{12} & Q_{12} & U_{12} & Z_{18} & Q_{16} & U_{16} \\ \hline & \{\text{id}\} & S_2 & S_3 & \{\text{id}\} & S_2 & S_3 \end{array} \quad (104)$$

Now we compare $\text{Stab}_{G_w}(f_0)$. It is sufficient to find enough elements so that the resulting group has half as many elements as $G_{\mathbb{Z}}$.

The cases Z_{12} and Z_{18} : Then

$$\varphi_1 : (x, y) \mapsto (e^{2\pi i w_x} x, e^{2\pi i w_y} y) \quad \text{satisfies} \quad (\varphi_1)_{\text{hom}} = M_h. \quad (105)$$

This is already sufficient. Here $G^{smar} = G_{\mathcal{R}}^{smar} = \{M_h^k \mid k \in \mathbb{Z}\}$.

The cases $Q_{12}, Q_{16}, U_{12}, U_{16}$: Here it is convenient to make use of the decomposition of the singularity f_0 into a sum of an A_l singularity g_0 in one variable and a D_{2m} singularity h_0 in two variables. In all four cases the weight system w' of the A_l singularity and the weight system

w'' of the D_{2m} singularity have denominators $l + 1$ and $2m - 1$ with $\gcd(l + 1, 2m - 1) = 1$. Therefore

$$\begin{aligned} \text{Stab}_{G_w}(f_0) &= \text{Stab}_{G_{w'}}(g_0) \times \text{Stab}_{G_{w''}}(h_0) \\ &\cong \mathbb{Z}_{l+1} \times \begin{cases} \mathbb{Z}_{2m-1} \times S_2 & \text{if } m \geq 3 \\ \mathbb{Z}_3 \times S_3 & \text{if } m = 2. \end{cases} \end{aligned} \quad (106)$$

In all four cases this group has half as many elements as $G_{\mathbb{Z}}$. $\square$

8. MORE ON $G_{\mathbb{Z}}$ FOR THE SIMPLE ELLIPTIC SINGULARITIES

This section is motivated by the paper [MS] of Milanov and Shen. They consider the 1-parameter families

$$x^p + y^q + z^r + t \cdot xyz, \quad t \in \Sigma \subset \mathbb{C} \quad (107)$$

of the simple elliptic singularities T_{pqr} with $(p, q, r) \in \{(3, 3, 3), (4, 4, 2), (6, 3, 2)\}$ which had also been used above in section 6. Here $\Sigma \subset \mathbb{C}$ is the complement of the finite set of parameters where the function in (107) has a non-isolated singularity. Remark that now $\chi := \text{lcm}(p, q, r) = p$.

In [MS] the groups of the transversal monodromies of these three families are studied, more precisely, the natural representations

$$\begin{aligned} \rho : \pi_1(\Sigma) &\rightarrow G_{\mathbb{Z}}, \\ \rho_1 : \pi_1(\Sigma) &\rightarrow \text{Aut}(Ml(f)_{1, \mathbb{Z}}, L), \\ \rho_{\neq 1} : \pi_1(\Sigma) &\rightarrow \text{Aut}(Ml(f)_{\neq 1, \mathbb{Z}}, L), \\ \bar{\rho}_{\neq 1} : \pi_1(\Sigma) &\rightarrow \text{Aut}(Ml(f)_{\neq 1, \mathbb{Z}}, L) / \langle M_h \rangle. \end{aligned} \quad (108)$$

By explicit computations they show

$$\ker(\rho_1) \subset \ker(\bar{\rho}_{\neq 1}). \quad (109)$$

They ask about a conceptual explanation of (109) and whether this might be true for other normal forms, i.e. other natural 1-parameter families of the simple elliptic singularities. Because of (109), there is an induced representation

$$\rho_W : \text{Im}(\rho_1) \rightarrow \text{Aut}(Ml(f)_{\neq 1, \mathbb{Z}}, L) / \langle M_h \rangle. \quad (110)$$

Then $\ker(\rho_W) \subset \text{Im}(\rho_1) \subset \text{Aut}(Ml(f)_{1, \mathbb{Z}}, L)$. One main result of [MS] is the following.

Theorem 8.1. *In all three cases, under an isomorphism $\text{Aut}(Ml(f)_{1, \mathbb{Z}}, L) \cong SL(2, \mathbb{Z})$ as in (18), the subgroup $\ker(\rho_W)$ is isomorphic to the principal congruence subgroup $\Gamma(p)$.*

Here $\Gamma(N) := \{A \in SL(2, \mathbb{Z}) \mid A \equiv \mathbf{1}_2 \pmod{N}\}$ (not $A \equiv \pm \mathbf{1}_2 \pmod{N}$) is the principal congruence subgroup of level N of $SL(2, \mathbb{Z})$.

In the following, we give results which complement (109) and theorem 8.1. We consider not a special 1-parameter family, but the biggest possible family of quasihomogeneous simple elliptic singularities. Define $(w_x, w_y, w_z) := (\frac{1}{p}, \frac{1}{q}, \frac{1}{r})$, define for any monomial its weighted degree $\deg_w(x^\alpha y^\beta z^\gamma) := \alpha w_x + \beta w_y + \gamma w_z$, and define

$$\mathbb{C}[x, y, z]_1 := \langle x^\alpha y^\beta z^\gamma \mid \deg_w(x^\alpha y^\beta z^\gamma) = 1 \rangle_{\mathbb{C}}, \quad (111)$$

$$R := \{f \in \mathbb{C}[x, y, z]_1 \mid f \text{ has an isolated singularity at } 0\}.$$

R is the complement of a hypersurface in the vector space $\mathbb{C}[x, y, z]_1$. The transversal monodromies of the family of singularities parametrized by R give the natural representation σ , the other representations are induced,

$$\begin{aligned} \sigma : \pi_1(R) &\rightarrow G_{\mathbb{Z}}, \\ \sigma_1 : \pi_1(R) &\rightarrow \text{Aut}(Ml(f)_{1, \mathbb{Z}}, L), \\ \sigma_{\neq 1} : \pi_1(R) &\rightarrow \text{Aut}(Ml(f)_{\neq 1, \mathbb{Z}}, L), \\ \bar{\sigma}_{\neq 1} : \pi_1(R) &\rightarrow \text{Aut}(Ml(f)_{\neq 1, \mathbb{Z}}, L) / \langle M_h \rangle. \end{aligned} \quad (112)$$

By the definition of G^{smar} in [He7, definition 3.1], $G^{smar} = \text{Im}(\sigma)$. Theorem 6.1 tells that the monodromy group $\text{Im}(\sigma)$ is as large as possible (up to $\pm \text{id}$ in the case T_{333}).

Theorem 8.2. *$\text{Im}(\sigma) = G_{\mathbb{Z}}$ in the cases T_{442} and T_{632} , and $G_{\mathbb{Z}} = \text{Im}(\sigma) \times \{\pm \text{id}\}$ in the case T_{333} (here it is important that the surface singularities are considered).*

The explicit information on $G_{\mathbb{Z}}$ in theorem 3.1 allows the following conclusion.

Corollary 8.3. *The analogue $\ker(\sigma_1) \subset \ker(\bar{\sigma}_{\neq 1})$ of (109) does not hold in the cases T_{333} and T_{442} . It holds in the case T_{632} , and there the analogue of (109) holds for any μ -constant family.*

Proof: By theorem 3.1 (c),

$$\begin{aligned} &\{g \in G_{\mathbb{Z}} \mid g|_{Ml(f)_1} = \text{id}\} \\ &= \{\text{id}|_{Ml(f)_1} \times (M_h|_{Ml_{\mathbb{C}}^{(1)}})^{\alpha} \times (M_h|_{Ml_{\mathbb{C}}^{(2)}})^{\beta} \times (M_h|_{Ml_{\mathbb{C}}^{(3)}})^{\gamma} \mid \\ &\quad (\alpha, \beta, \gamma) \in \mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r \text{ with } \frac{\alpha}{p} + \frac{\beta}{q} + \frac{\gamma}{r} \equiv 0 \pmod{1}\} \times U_2. \end{aligned}$$

In the cases T_{442} and T_{632} this is isomorphic to $\ker(\sigma_1)/\ker(\sigma)$, in the case T_{333} $\ker(\sigma_1)/\ker(\sigma)$ is isomorphic to this group or a subgroup of

this group of index 2. In the cases T_{442} and T_{333} , already the factor U_2 in this group is an obstruction to the analogue of (109).

In the case T_{632} , U_2 is trivial and

$$\{g \in G_{\mathbb{Z}} \mid g|_{Ml(f)_1} = \text{id}\} = \{M_h^\alpha \mid \alpha \in \mathbb{Z}\} \quad (113)$$

(because above α determines β and γ uniquely in the case $(p, q, r) = (6, 3, 2)$). Therefore the analogue of (109) holds in the case T_{632} for the family parametrized by R and for any subfamily. $\square$

(109) is used in [MS] in order to define ρ_W and the group $\ker(\rho_W)$. But because $M_h|_{Ml(f)_1} = \text{id}$, it is obvious that the group $\ker(\rho_W)$ coincides with

$$\{g|_{Ml(f)_{1,\mathbb{Z}}} \mid g \in \text{Im}(\sigma), g|_{Ml(f)_{\neq 1,\mathbb{Z}}} = \text{id}\} \subset \text{Aut}(Ml(f)_{1,\mathbb{Z}}, L). \quad (114)$$

And the analogue of this group can be defined for any μ -constant family, whether or not it satisfies the analogue of (109). Our main result in this section is theorem 8.4. Our proof uses theorem 3.1. A different proof of theorem 8.4 was given by Kluitmann in [Kl, III 2.4 Satz, page 66]. Theorem 8.4 shows that the group $\Gamma(p)$ turns up naturally within the maximal possible μ -constant family, which is parametrized by R , and it shows the part $\ker(\rho_W) \subset \Gamma(p)$ of the equality $\ker(\rho_W) = \Gamma(p)$ in theorem 8.1.

Theorem 8.4. *In all three cases, under an isomorphism $\text{Aut}(Ml(f)_{1,\mathbb{Z}}, L) \cong SL(2, \mathbb{Z})$ as in (18), the subgroup*

$$\{g|_{Ml(f)_{1,\mathbb{Z}}} \mid g \in G_{\mathbb{Z}}, g|_{Ml(f)_{\neq 1,\mathbb{Z}}} = \text{id}\} \subset \text{Aut}(Ml(f)_{1,\mathbb{Z}}, L).$$

is isomorphic to the principal congruence subgroup $\Gamma(p)$.

Proof: We use the notations and objects in the proof of theorem 3.1. $Ml_{\mathbb{C}}^{(1)}$ was defined in (31). Define

$$Ml_{\mathbb{Z}}^{(1)} := Ml_{\mathbb{C}}^{(1)} \cap Ml(f), \quad (115)$$

and analogously $Ml_{\mathbb{Z}}^{(2)}$ and $Ml_{\mathbb{Z}}^{(3)}$. Then

$$Ml_{\mathbb{Z}}^{(1)} = (M_h - \text{id})(Ml_{\mathbb{Z}}^{[1]}).$$

Because of (32), this image is generated as a $\mathbb{Z}$ -lattice by

$$\delta_2 - (\delta_1 + \tilde{b}_1), \delta_3 - \delta_2, \dots, \delta_{p-1} - \delta_{p-2}, -(\delta_1 + \dots + \delta_{p-1}) - \delta_{p-1}, \quad (116)$$

respectively by

$$\delta_1 + (p-1)\delta_2, \delta_3 - \delta_2, \dots, \delta_{p-1} - \delta_{p-2}, p\delta_2 - \delta_{\mu-1} + \delta_{\mu}. \quad (117)$$

$Ml_{\mathbb{Z}}^{(2)}$ and $Ml_{\mathbb{Z}}^{(3)}$ (if $r \geq 3$) are generated by the analogous elements. If $r = 2$ then

$$Ml_{\mathbb{Z}}^{(3)} = \mathbb{Z} \cdot (2\delta_{\mu-2} - \delta_{\mu-1} + \delta_{\mu}). \quad (118)$$

In any case, the sum $Ml_{\mathbb{Z}}^{(1)} \oplus Ml_{\mathbb{Z}}^{(2)} \oplus Ml_{\mathbb{Z}}^{(3)}$ is a sublattice of finite index of the primitive sublattice $Ml(f)_{\neq 1, \mathbb{Z}}$ in $Ml(f)$. Observe that $q|p$ and $r|p$ in all three cases. The lattice $Ml(f)_{\neq 1, \mathbb{Z}}$ is generated by

$$\begin{aligned} & \delta_1 + (p-1)\delta_2, \delta_3 - \delta_2, \dots, \delta_{p-1} - \delta_{p-2}, p\delta_2 - \delta_{\mu-1} + \delta_{\mu}, \quad (119) \\ & \delta_p + (q-1)\delta_{p+1}, \delta_{p+2} - \delta_{p+1}, \dots, \delta_{p+q-2} - \delta_{p+q-3}, \frac{p}{q}\delta_2 - \delta_{p+1}, \\ & \delta_{p+q-1} + (r-1)\delta_{p+q}, \delta_{p+q+1} - \delta_{p+q}, \dots, \delta_{\mu-2} - \delta_{\mu-3}, \frac{p}{r}\delta_2 - \delta_{p+q}, \end{aligned}$$

if $r \geq 3$. If $r = 2$ then the third line has to be replaced by

$$\frac{p}{r}\delta_2 - \delta_{\mu-2}.$$

In any case, one sees

$$Ml(f) = Ml(f)_{\neq 1, \mathbb{Z}} \oplus \mathbb{Z} \cdot \delta_2 \oplus \mathbb{Z} \cdot \delta_{\mu}. \quad (120)$$

One also calculates

$$\tilde{b}_1 = \delta_{\mu-1} - \delta_{\mu} = \gamma_1 + p\delta_2 \quad (121)$$

$$\text{with } \gamma_1 := -(p\delta_2 - \delta_{\mu-1} + \delta_{\mu}) \in Ml(f)_{\neq 1, \mathbb{Z}},$$

$$\tilde{b}_2 = \gamma_2 + p\delta_{\mu}, \quad (122)$$

$$\begin{aligned} \text{with } \gamma_2 := & \sum_{i=1}^{p-1} (p-i)\delta_i + \sum_{i=1}^{q-1} \frac{p}{q}(q-i)\delta_{p-1+i} \\ & + \sum_{i=1}^{r-1} \frac{p}{r}(r-i)\delta_{p+q-2+i} + p\delta_{\mu-1} - p\delta_{\mu} \in Ml(f)_{\neq 1, \mathbb{Z}}. \end{aligned}$$

One sees also

$$Ml(f) \cap (\mathbb{Q} \cdot \gamma_1 + \mathbb{Q} \cdot \gamma_2) = \mathbb{Z} \cdot \gamma_1 \oplus \mathbb{Z} \cdot \gamma_2. \quad (123)$$

For any matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ define the automorphism $f : Ml(f)_{\mathbb{Q}} \rightarrow Ml(f)_{\mathbb{Q}}$ by

$$f(\tilde{b}_1) := a\tilde{b}_1 + c\tilde{b}_2, \quad f(\tilde{b}_2) := b\tilde{b}_1 + d\tilde{b}_2, \quad f|_{Ml(f)_{\neq 1, \mathbb{Z}}} := \text{id}. \quad (124)$$

It respects L because the decomposition $Ml(f)_{\mathbb{Q}} = Ml(f)_{1, \mathbb{Q}} \oplus Ml(f)_{\neq 1, \mathbb{Q}}$ is left and right orthogonal with respect to L . It is an

automorphism of $Ml(f)$ if and only if $f(\delta_2)$ and $f(\delta_\mu)$ are in $Ml(f)$. One calculates

$$\begin{aligned} f(\delta_2) &= a\delta_2 + c\delta_\mu + \frac{1}{p}(a-1)\gamma_1 + \frac{1}{p}c\gamma_2, \\ f(\delta_\mu) &= b\delta_2 + d\delta_\mu + \frac{1}{p}b\gamma_1 + \frac{1}{p}(d-1)\gamma_2. \end{aligned}$$

In view of (123) this shows

$$f \in G_{\mathbb{Z}} \iff \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \mathbf{1}_2 \pmod{p} \stackrel{\text{def.}}{\iff} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma(p). \quad \square$$

REFERENCES

- [AGV1] V.I. Arnold, S.M. Gusein-Zade, A.N. Varchenko: Singularities of differentiable maps, volume I. Birkhäuser, Boston, Basel, Stuttgart, 1985.
- [AGV2] V.I. Arnold, S.M. Gusein-Zade, A.N. Varchenko: Singularities of differentiable maps, volume II. Birkhäuser, Boston 1988.
- [Eb1] W. Ebeling: Milnor lattices and geometric bases of some special singularities. In: Noeuds, tresses et singularités, Monographie No **31** de l'enseignement Mathématique, Genève, 1983, 129–146.
- [Eb2] W. Ebeling: Functions of several complex variables and their singularities. Graduate Studies in Mathematics **83**, AMS, 2007.
- [Ga] A.M. Gabrielov: Dynkin diagrams of unimodal singularities. *Funct. Anal. Appl.* **8** (1974), 192–196.
- [He1] C. Hertling: Analytische Invarianten bei den unimodularen und bimodularen Hyperflächensingularitäten. Doctoral thesis. Bonner Math. Schriften **250**, Bonn 1992.
- [He2] C. Hertling: Ein Torellisatz für die unimodalen und bimodularen Hyperflächensingularitäten. *Math. Ann.* **302** (1995), 359–394.
- [He3] C. Hertling: Brieskorn lattices and Torelli type theorems for cubics in $\mathbb{P}^3$ and for Brieskorn-Pham singularities with coprime exponents. In: Singularities, the Brieskorn anniversary volume. *Progress in Mathematics* **162**. Birkhäuser Verlag, Basel-Boston-Berlin 1998, pp. 167–194.
- [He4] C. Hertling: Classifying spaces and moduli spaces for polarized mixed Hodge structures and for Brieskorn lattices. *Compositio Math.* **116** (1999), 1–37.
- [He5] C. Hertling: Generic Torelli for semiquasihomogeneous singularities. In: Trends in singularities (eds. A. Libgober, M. Tibar), Birkhäuser Verlag, Basel 2002, 115–140.
- [He6] C. Hertling: Frobenius manifolds and moduli spaces for singularities. *Cambridge Tracts in Mathematics* **151**, Cambridge University Press, 2002.
- [He7] C. Hertling: μ -constant monodromy groups and marked singularities. *Ann. Inst. Fourier, Grenoble* **61.7** (2011), 2643–2680.
- [Ka] R. Kaenders: The Seifert form of a plane curve singularity determines its intersection multiplicities. *Indag. Mathem. N.S.* **7.2** (1996), 185–197.
- [Kl] P. Kluitmann: Ausgezeichnete Basen erweiterter affiner Wurzelgitter. Doctoral thesis. Bonner Math. Schriften **185**, Bonn 1987.

- [Ku] Va.S. Kulikov: Mixed Hodge structures and singularities. Cambridge tracts in mathematics **132**, Cambridge University Press, 1998.
- [LR] Lê Dũng Tráng, C.P. Ramanujam: The invariance of Milnor's number implies the invariance of the topological type. Amer. J. of Math. **98** (1973), 67–78.
- [MW] F. Michel, C. Weber: Sur le rôle de la monodromie entière dans la topologie des singularités. Ann. Inst. Fourier Grenoble **36** (1986), 183–218.
- [MS] T. Milanov, Y. Shen: The modular group for the total ancestor potential of Fermat simple elliptic singularities. Comm. Number Theor. Phys. **8** (2014), 329–368.
- [Mi] J. Milnor: Singular points of complex hypersurfaces. Ann. Math. Stud. vol. 61, Princeton University Press 1968.
- [Or] P. Orlik: On the homology of weighted homogeneous polynomials. In: Lecture Notes in Math. 298, Springer, Berlin, 1972.
- [SaK] K. Saito: Einfach-elliptische Singularitäten. Invent. Math. **23** (1974), 289–325.
- [SaM] M. Saito: Period mapping via Brieskorn modules. Bull. Soc. math. France **119** (1991), 141–171.

FALKO GAUSS, UNIVERSITÄT MANNHEIM, LEHRSTUHL FÜR MATHEMATIK VI,
SEMINARGEBAUDE A 5, 6, 68131 MANNHEIM, GERMANY
E-mail address: `gauss@math.uni-mannheim.de`

CLAUS HERTLING, UNIVERSITÄT MANNHEIM, LEHRSTUHL FÜR MATHEMATIK VI,
SEMINARGEBAUDE A 5, 6, 68131 MANNHEIM, GERMANY
E-mail address: `hertling@math.uni-mannheim.de`